

## DIETARY TURMERIC (*Curcuma longa*) ENHANCES GROWTH, ANTIOXIDANT DEFENSE, AND IMMUNITY IN *Cirrhinus mrigala*

Ali Hassan<sup>1</sup>, Kajeen Hassan Jasim<sup>2</sup>, Ayesha Shahid<sup>3</sup>, Nizar J. Hussein<sup>4</sup>, Hafiza Aqsa Azam<sup>5</sup>, and Muhammad Owais<sup>1,\*</sup>

<sup>1</sup>Department of Zoology, Ghazi University, Dera Ghazi Khan, Pakistan.

<sup>2</sup>Medical Laboratory Department, College of Health Sciences, Cihan University-Dohuk, Kurdistan Region, Iraq.

<sup>3</sup>Department of Zoology, Government College University Lahore, Pakistan.

<sup>4</sup>Environmental Sciences Department, College of Science, University of Zakho, Kurdistan Region of Iraq.

<sup>5</sup>Department of Zoology, Government College University, Faisalabad, Pakistan.

\* Corresponding author email: [owaisgulmuhammad@gmail.com](mailto:owaisgulmuhammad@gmail.com)

Received: 09Jul 2025 Accepted:31 Jul 2025 Published:01 Oct 2025 <https://doi.org/10.25271/sjuoz.2025.13.4.1688>

### ABSTRACT:

Aquaculture is pivotal in addressing global food security, yet intensive practices compromise fish health and productivity. This study investigated the efficacy of dietary turmeric (*Curcuma longa*) supplementation as a natural alternative in *Cirrhinus mrigala*, a species critical to South Asian aquaculture. A 60-day feeding trial evaluated four diets: control (0% turmeric), 0.5%, 1.0%, and 1.5% turmeric inclusion. Results demonstrated that 1.0% turmeric significantly enhanced growth performance, with a 28% increase in final body weight ( $19.6 \pm 1.4$  g vs.  $15.7 \pm 1.1$  g in control;  $p < 0.01$ ) and improved feed conversion ratio ( $1.4 \pm 0.2$  vs.  $1.7 \pm 0.2$ ;  $p = 0.004$ ). Antioxidant enzyme activities (SOD, CAT, GPx) increased by 34–53%, while lipid peroxidation (MDA) decreased by 29% ( $p = 0.001$ ), indicating robust oxidative stress mitigation. Immunological assays revealed a 65% rise in lysozyme activity and 70% higher phagocytic capacity ( $p < 0.01$ ), underscoring turmeric's immunostimulatory potential. The hepatosomatic index decreased by ( $1.3 \pm 0.2\%$  vs.  $1.4 \pm 0.2\%$ ;  $p = 0.038$ ), suggesting increased metabolic efficiency. These findings highlight 1.0% turmeric as the optimal dosage, offering a sustainable strategy to augment aquaculture productivity while reducing reliance on antibiotics.

**KEYWORDS:** *Curcuma longa*, Growth, Antioxidant Defense, Immunity, *Cirrhinus mrigala*

### 1. INTRODUCTION

Fish farming has emerged as a vital approach to meet the growing global demand for seafood, currently supplying more than 50% of the fish eaten worldwide (Khanjani *et al.*, 2024; Nabi *et al.*, 2025). The growth of aquaculture is crucial for developing economies, as seafood provides a cost-effective, nutritious protein source (Munguti *et al.*, 2024; Rashaia *et al.*, 2025). Large-scale aquaculture has made significant advancements due to improvements in health and sustainability in potentially different environmental concerns, while chronic stress and high stocking densities decrease the farm productivity and fish welfare (Hematyar *et al.*, 2024; Yadav *et al.*, 2024). The use of antibiotics has become a major global health issue in aquaculture because many fish farms rely on antibiotics, and 80% antibiotic resistance genes were found in aquaculture water samples of different fish farms (Hossain *et al.*, 2022; Rasul *et al.*, 2025). To minimise these emerging current issues, the global regulatory authorities restricted the use of antibiotics in aquaculture and also addressed the alternative health solutions, such as plant-based feed additives, which emerged as a promising alternative that can give numerous health benefits (Mithuna *et al.*, 2024; Pitiot *et al.*, 2025; Rasheed *et al.*, 2024).

*Cirrhinus mrigala* (freshwater carp) is an important aquaculture species found in South Asia, valued for its nutritional benefits, delicious taste, and fast growth (Meitei *et al.*, 2025). *C. mrigala* plays a vigorous role in food security and supports the various small-scale fish farmers of different regions of South Asia (Bhattacharya *et al.*, 2024; Debnath *et al.*, 2025). Moreover, during the intensive farming of *C. mrigala*, especially during the monsoon seasons, it becomes a greater chance of bacterial infections that cause stunted growth in unhealthy water conditions, and these environmental changes disturb the fish's immunity and disrupt the disease patterns (Al Sulivany *et al.*, 2024; Kasihmuddin *et al.*, 2024).

*Curcuma longa* L., commonly known as turmeric, contains bioactive compounds such as bisdemethoxycurcumin and demethoxycurcumin (Hafeez *et al.*, 2024; Roney *et al.*, 2025). It is a powerful natural feed supplement that maintains unhealthy water conditions and offers significant health and growth benefits (Bhatt *et al.*, 2024; Fan *et al.*, 2025). Moreover, it is efficiently absorbed by fish through commercial feed (Abdelkarim *et al.*, 2025; Naghdi *et al.*, 2024). Turmeric also plays a powerful role as an antioxidant, boosting the fish's natural defense systems and neutralizing harmful free radicals through various biological pathways (Bouyanya *et al.*, 2024; Esmaelzadeh *et al.*, 2024). In aquaculture, it improved the biosynthesis of antioxidant

\* Corresponding author

This is an open access under a CC BY-NC-SA 4.0 license (<https://creativecommons.org/licenses/by-nc-sa/4.0/>)

enzymes, including Superoxide dismutase and catalase, and reduced cellular oxidative stress markers (Fan *et al.*, 2025; Jomova *et al.*, 2024).

*C. longa* supplementation in aquaculture enhances various physiological adaptations, such as significant gastrointestinal enhancements, improved intestinal villi architecture, and enterocyte regeneration in specimens receiving dietary turmeric inclusion (Khaldoun *et al.*, 2025; Song *et al.*, 2024).

Due to wide research on the aquacultural benefits of *C. longa* the optimum application procedures for *C. mrigala* cultivation require further explanation; However, turmeric's effective dosage typically differs, typically ranging from 0.5-2% of feed (Komal *et al.*, 2024; Wildhaber *et al.*, 2023). The research gaps remain regarding *C. longa* long-term effects on meat quality and fish health (Ringø *et al.*, 2023). This experimental study investigates the dietary turmeric supplementation in the diet of *C. mrigala* to check the growth, antioxidant defence, and immunity.

## 2. MATERIALS AND METHODS

The current experimental study was conducted at the Saline Water Aquaculture Research Centre (SWARC), Muzaffargarh. The 216 juvenile experimental fish (*C. mrigala*) with an average weight of  $5.3 \pm 0.14$  g were purchased from the Fish Hatchery, SWARC, Muzaffargarh. Four treatment groups with three replicates were designed: T1 (control, 0% turmeric), T2 (0.5% turmeric), T3 (1.0% turmeric), and T4 (1.5% turmeric). Before the experiment, fish underwent a 15-day acclimation period in a 2000L fiberglass tank with continuous aeration, daily 30% water exchange, and twice-daily feeding of a standard 32% protein diet at 3% body weight. Healthy fish showing active swimming

behavior, clear eyes, intact fins, uniform coloration, and bright red gills were selected and randomly stocked into twelve 300L experimental tanks (18 fish per tank). Each tank was equipped with individual water circulation, separate aeration, 1mm mesh covers, and waste collection trays. Water quality was rigorously maintained at 28-30°C (using submersible heaters), dissolved oxygen  $>5.8$  mg/L, pH 7.4-7.8 (adjusted with sodium bicarbonate), and ammonia  $<0.1$  mg/L. All parameters were monitored three times daily using calibrated Hanna HI98194 meters to ensure experimental consistency throughout the 60-day trial period. The research framework enabled precise assessment of *C. longa*'s physiological impacts on *C. mrigala* under standardized aquaculture parameters (Hoseini & Al Sulivany, 2024; Owais *et al.*, 2024).

### Experimental Diets and Feeding Protocol:

Four isonitrogenous (32% crude protein) experimental feeds were formulated with graded *C. longa* inclusions (0-1.5%). All diets utilized premium fishmeal and soybean meal protein sources with complete vitamin-mineral fortification. Feed production required accurate ingredient measurement, homogenization, and extrusion into 2mm pellets using 35°C purified water. After 48 hours of drying at 28°C, pellets were vacuum-packed and frozen to maintain nutritional quality. The 60-day experiment involved twice-daily feedings (3% biomass), with rations adjusted weekly. Uneaten pellets were removed after 45 minutes to prevent water contamination. The study employed twelve 300-litre tanks (18 fish/tank), randomly assigned to four treatment groups with three replicates (Owais *et al.*, 2023; Al Sulivany *et al.*, 2024a; Abdulrahman & Al Sulivany, 2025).

**Table 1:** Composition of experimental diets supplemented with graded levels of turmeric powder (g/kg)

Ingredients	T1 (0%)	T2 (0.5%)	T3 (1.0%)	T4 (1.5%)
Fish meal	190	190	190	190
Soybean meal	80	80	80	80
Cottonseed cake	110	110	110	110
Wheat bran	310	308.5	307	305.5
Wheat pollard	130	130	130	130
Maize germ	100	100	100	100
Soybean oil	18	18	18	18
Vitamin-mineral premix*	12	12	12	12
Turmeric powder	0	1.5	3.0	4.5
Total (g)	950	950	950	950

Note: \*Vitamin-mineral premix per kg of feed: Vitamin A – 7200 IU, Vitamin D<sub>3</sub> – 1600 IU, Vitamin E – 200 IU, Vitamin C – 90 mg, Vitamin B<sub>1</sub> – 1.5 mg, Vitamin B<sub>2</sub> – 10 mg, Vitamin B<sub>6</sub> – 8 mg, Vitamin B<sub>12</sub> – 0.02 mg, Niacin – 55 mg, Pantothenic acid – 40 mg, Folic acid – 0.08 mg, Biotin – 1.8 mg, Copper – 6 mg, Iron – 110 mg, Zinc – 30 mg, Manganese – 18 mg, Iodine – 0.6 mg, Selenium – 0.2 mg, and Cobalt – 0.12 mg.

### Growth Performance Assessment:

Following the 60-day feeding period, all *C. mrigala* specimens underwent a 24-hour fasting period to ensure complete gut clearance before final measurements. Individual fish were carefully weighed using a precision digital balance (accuracy  $\pm 0.01$ g) to determine final body weights (Al Sulivany *et al.*, 2024b).

$$\text{Weight gain (g)} = \text{Final weight} - \text{Initial weight}$$

$$\text{Specific growth rate (\%/day)} = [(\ln \text{final weight} - \ln \text{initial weight})/\text{days}] \times 100$$

$$\text{Feed conversion ratio} = \text{Total feed intake (g)}/\text{Weight gain (g)}$$

$$\text{Protein efficiency ratio} = \text{Weight gain (g)}/\text{Protein intake (g)}$$

$$\text{Survival rate (\%)} = (\text{Final fish number}/\text{Initial fish number}) \times 100$$

Hepatosomatic index (\%) = (Liver weight/Body weight)  $\times 100$   
Measurements followed standard aquaculture protocols using calibrated equipment. Three technicians independently verified all calculations to ensure accuracy. The hepatosomatic index was determined through careful dissection and weighing of liver tissues from sampled specimens.

### Antioxidant Enzyme Activities:

To evaluate antioxidant and immune responses, blood samples were obtained from sedated fish following standard protocols. For each treatment group, blood was collected from 3 fish per tank, for a total of 9 fish per treatment group (n = 9).

Superoxide dismutase (SOD) activity was assessed by measuring pyrogallol oxidation at 560 nm, while catalase (CAT) activity was determined by observing hydrogen peroxide breakdown at 240 nm. Glutathione peroxidase (GPx) activity was assessed by measuring NADPH depletion at 340 nm. Oxidative damage was evaluated by determining malondialdehyde (MDA) levels at 532 nm via thiobarbituric acid assay. All enzyme levels were normalized to total protein concentration, determined using the Bradford assay (595 nm). Adapted for fish tissue analysis, measurements were performed in triplicate with strict quality checks, following standardized protocols (Taysi *et al.*, 2024; Verdi *et al.*, 2024).

#### Innate Immune Parameters:

Serum lysozyme levels were assessed by measuring the lysis of *Micrococcus lysodeikticus*. A mixture of 25  $\mu$ L serum and 175  $\mu$ L bacterial suspension (0.2 mg/mL PBS) was prepared, and absorbance at 450 nm was monitored for 5 minutes. Lysozyme activity units were calculated based on a 0.001 minute $^{-1}$  absorbance decline at 450 nm. Head kidney leukocytes were separated by Percoll gradient centrifugation to distinct feasible phagocytic cells. The purified cells were then incubated with fluorescently labelled bacteria at 25 °C for 30–60 minutes to measure phagocytic activity. After incubation, non-internalized particles were washed off, and the cells were examined using flow cytometry. *cerevisiae* (10% v/v) for 60 minutes at 28°C. Following fixation and staining, phagocytic capacity was quantified as the percentage of cell-positive leukocytes per 100 microscopic fields. All assays were conducted in triplicate with matched controls under aseptic conditions to prevent microbial contamination (Díaz-Puertas *et al.*, 2024; Elkhaiat *et al.*, 2025; Lulijwa *et al.*, 2019).

#### Statistical Analysis:

Data were validated through comprehensive statistical testing. Distribution normality was assessed with Shapiro-Wilk ( $\alpha=0.05$ ), and variance uniformity was evaluated via Levene's test. Treatment differences were analyzed by one-way ANOVA, with Tukey's HSD post-hoc tests applied for significant results ( $p<0.05$ ). Analyses used IBM SPSS v26.0 (Armonk, NY), with a significance threshold at  $p<0.05$ . Values represent mean  $\pm$  SE ( $n=3$  biological replicates/treatment).

### 3. RESULTS

The experimental investigation evaluating graded levels of dietary turmeric *C. longa* supplementation in *C. mrigala* revealed significant improvements across multiple physiological parameters. The 1.0% turmeric inclusion consistently demonstrated optimal efficacy, establishing a clear dose-response relationship.

Final body weights showed progressive enhancement with increasing turmeric concentrations. The average initial weight was  $15.3 \pm 1.1$  g, while the 0.5%, 1.0%, and 1.5% supplemented groups reached  $17.4 \pm 1.2$  g ( $p < 0.05$ ),  $19.6 \pm 1.4$  g ( $p < 0.01$ ), and  $17.8 \pm 1.2$  g ( $p < 0.003$ ), respectively. Weight gain followed an identical pattern, with the 1.0% group achieving  $14.3 \pm 1.3$  g compared to control values of  $10.4 \pm 0.9$  g ( $p = 0.002$ ). SGR exhibited maximum improvement in the 1.0% treatment ( $2.11 \pm 0.06\%$ /day) versus controls ( $1.72 \pm 0.05\%$ /day;  $p = 0.001$ ). FCR demonstrated similar optimization, decreasing from  $1.7 \pm 0.2$  in controls to  $1.4 \pm 0.2$  in the 1.0% group ( $p = 0.004$ ). PER increased from  $2.2 \pm 0.2$  to  $2.7 \pm 0.3$  ( $p = 0.005$ ) at the optimal supplementation level. Survival rates remained consistently high across all treatments (92–98%), showing no statistically significant variation ( $p = 0.213$ ). The hepatosomatic index (HSI) displayed a modest but significant reduction in the 1.0% group ( $1.3 \pm 0.2\%$ ) compared to control ( $1.4 \pm 0.2\%$ ;  $p = 0.038$ ), potentially indicating improved hepatic metabolic efficiency (Table 2).

**Table 2:** Effects of Dietary Turmeric Supplementation on Weight Gain, Feed Efficiency, and Hepatosomatic Index in *Cirrhinus mrigala*

Parameter	Control	0.5% Turmeric	1.0% Turmeric	1.5% Turmeric	p-value
<b>Initial Weight (g)</b>	$5.3 \pm 0.14$	$5.3 \pm 0.12$	$5.3 \pm 0.16$	$5.3 \pm 0.13$	0.06
<b>Final Weight (g)</b>	$15.7 \pm 1.1$	$17.4 \pm 1.2^*$	$19.6 \pm 1.4^{**}$	$17.8 \pm 1.2^*$	0.003
<b>Weight Gain (g)</b>	$10.4 \pm 0.9$	$12.1 \pm 1.1^*$	$14.3 \pm 1.3^{**}$	$12.5 \pm 1.0^*$	0.002
<b>SGR (%/day)</b>	$1.72 \pm 0.05$	$1.94 \pm 0.06^*$	$2.11 \pm 0.06^{**}$	$2.04 \pm 0.08^*$	0.001
<b>FCR</b>	$1.7 \pm 0.2$	$1.5 \pm 0.3^*$	$1.4 \pm 0.2^{**}$	$1.5 \pm 0.2^*$	0.004
<b>PER</b>	$2.2 \pm 0.2$	$2.43 \pm 0.3^*$	$2.7 \pm 0.4^{**}$	$2.5 \pm 0.3^*$	0.005
<b>Survival Rate (%)</b>	$92 \pm 3$	$97 \pm 3$	$98 \pm 4$	$95 \pm 5$	0.213
<b>HSI (%)</b>	$1.4 \pm 0.2$	$1.4 \pm 0.4$	$1.3 \pm 0.2^*$	$1.4 \pm 0.1$	0.038

Note: Values represent mean  $\pm$  SD ( $n = 3$ ). Asterisks indicate significant differences compared to control: \* $p < 0.05$ , \*\* $p < 0.01$ . SGR = specific growth rate; FCR = feed conversion ratio; PER = protein efficiency ratio; HSI = hepatosomatic index. One-way ANOVA followed by Tukey's test was used for statistical analysis ( $p < 0.05$ ) \*

The antioxidant profile showed marked improvements with turmeric administration. SOD activity increased from  $13.2 \pm 1.4$  U/mg in controls to  $17.7 \pm 1.8$  U/mg in the 1.0% group ( $p = 0.003$ ). CAT activity rose from  $44.5 \pm 3.2$  U/mg to  $59.6 \pm 4.4$  U/mg ( $p = 0.008$ ), while (GPx) increased from  $17.3 \pm 2.2$  U/mg to  $26.4 \pm 2.7$  U/mg ( $p = 0.005$ ). These enzymatic enhancements corresponded with a 29% reduction in MDA levels ( $3.1 \pm 0.4$  nmol/mg to  $2.2 \pm 0.2$  nmol/mg;  $p = 0.001$ ), indicating significantly reduced lipid peroxidation.

Lysozyme activity, a crucial component of innate immunity, increased by 65% in the 1.0% group ( $1.7 \pm 0.2$  U/mL to  $2.8 \pm 0.3$  U/mL;  $p = 0.002$ ). Phagocytic capacity showed even greater enhancement, rising by 70% ( $21.5 \pm 2.7\%$  to  $36.6 \pm 3.8\%$ ;  $p = 0.006$ ). These immunological parameters demonstrated the strongest response at the 1.0% inclusion level, with slightly reduced efficacy at higher concentrations.

**Table 3:** Antioxidant enzyme activities and immune parameters of *C. mrigala* fed turmeric-supplemented diets for 60 days

Parameter	Control	0.5% Turmeric	1.0% Turmeric	1.5% Turmeric	p-value
<b>SOD (U/mg)</b>	$13.2 \pm 1.4$	$15.2 \pm 1.6^*$	$17.7 \pm 1.8^{**}$	$16.2 \pm 1.4^{**}$	0.003
<b>CAT (U/mg)</b>	$44.5 \pm 3.2$	$51.3 \pm 3.2^*$	$59.6 \pm 4.4^{**}$	$54.9 \pm 3.5^{**}$	0.008
<b>GPx (U/mg)</b>	$17.3 \pm 2.2$	$22.5 \pm 2.3^*$	$26.4 \pm 2.7^{**}$	$24.3 \pm 2.4^{**}$	0.005
<b>MDA (nmol/mg)</b>	$3.1 \pm 0.4$	$2.5 \pm 0.3^*$	$2.2 \pm 0.2^{**}$	$2.4 \pm 0.3^{**}$	0.001
<b>Lysozyme (U/mL)</b>	$1.7 \pm 0.2$	$2.4 \pm 0.3^*$	$2.8 \pm 0.3^{**}$	$2.3 \pm 0.4^{**}$	0.002
<b>Phagocytic activity (%)</b>	$21.5 \pm 2.7$	$27.4 \pm 3.2^*$	$36.6 \pm 3.8^{**}$	$32.2 \pm 3.4^{**}$	0.006

Note: Values are means  $\pm$  SD of three replicates. Significant differences compared to control group are indicated by \* ( $p < 0.05$ ) and \*\* ( $p < 0.01$ ). SOD = superoxide dismutase; CAT = catalase; GPx = glutathione peroxidase; MDA = malondialdehyde. Statistical analysis was performed using one-way ANOVA followed by Tukey's HSD test.

#### 4. DISCUSSION

This study examined the effect of *C. longa* 1.0% dietary supplementation in *C. mrigala*, which enhanced growth performance, increased antioxidant capacity, and improved immune responses. These results also align with the previous studies on turmeric's multifunctional advantages in aquaculture nutrition (Asad *et al.*, 2025; El-Gogary *et al.*, 2025; Khieokhajonhet *et al.*, 2023; Wei *et al.*, 2024), which describe that turmeric dietary supplementation in fish improved growth parameters viz, body weight, higher SGR, FCR, nutrient utilization, and metabolic efficiency. Similar growth performance was recorded in Nile tilapia with the dietary supplementation of curcumin through enhanced digestive function and intestinal health (Abdel-Ghany *et al.*, 2023; Kusi *et al.*, 2025; Sulivany *et al.*, 2024b). The present study describes the reduced HIS in the 1.0% turmeric that indicates enhanced hepatic efficiency. Similarly, Bhatt *et al.* (2024) also studied the reduced HSI in cyprinids.

The present study highlights the marked rise in antioxidant properties of critical defense enzymes

Viz: SOD, CAT, and GPx, and reduction in oxidative stress indicators like MDA. These results also align with the previous studies on turmeric, which also caused protective effects, and a marked rise in antioxidant properties was observed in common carp (Bin-Ammar *et al.*, 2024; Eissa *et al.*, 2024). Tahir *et al.* (2024) and Zheng *et al.* (2025) studied the turmeric antioxidant effects mechanism and explained that bioactive constituents directly scavenge reactive oxygen species, simultaneously activating the body's innate antioxidant defenses via the Nrf2

signaling pathway. Metwaly *et al.* (2025) also studied the Turmeric defensive properties in aquaculture systems, especially during ecological stressors such as high stocking densities and fluctuating water parameters that can lead to severe oxidative damage.

Turmeric supported immune function in the studied fish, as seen in improved infection-fighting enzyme levels and more efficient pathogen removal by immune cells, both crucial parts of fish immune defenses. These results align with Alsubaie *et al.* (2025) work showing curcumin-enriched feed increased disease resistance mechanisms in grouper fish. Turmeric boosts fish immunity by activating key defense pathways (NF- $\kappa$ B and MAPK), which increase protective protein production and disease resistance (Shin *et al.*, 2024; Wang *et al.*, 2025). These findings are especially valuable as fish farmers increasingly need natural alternatives to replace antibiotics and address drug resistance issues (Neil *et al.*, 2024).

Our study found the 1.0% turmeric dosage worked best, matching previous fish nutrition research. Komal *et al.* (2024) reported identical results in tilapia, where higher 1.5% doses became less effective - likely because the fish could only use so much of the active compounds. This dosage pattern appears across species, as Jasim *et al.* (2024) observed similar limits in chickens, where too much curcumin disrupted fat processing. While this study provides valuable insights, some questions remain unanswered. We did not examine how long-term turmeric use affects meat quality or breeding success in fish. Next steps should investigate these areas, along with testing turmeric combinations with other plant-based additives like garlic or thyme oil to create more effective feeds (Antache *et al.*, 2025; Khaldoun *et al.*, 2025; Rasheed *et al.*, 2024). Advanced genetic

analysis methods could also help pinpoint exactly how turmeric influences nutrient processing and immune system activation at the molecular level (Zou *et al.*, 2025).

## CONCLUSION

Our research shows that adding 1.0% turmeric to feed delivers the best results for boosting growth, fighting oxidative stress, and strengthening immunity in *C. mrigala*. These findings prove turmeric works effectively as a natural replacement for artificial additives in fish farming. While these results are promising, more studies should examine extended use effects and the most effective ways to administer turmeric. This work contributes to more sustainable fish production by enhancing welfare and yields naturally. Turmeric represents an eco-conscious option for aquaculture operations seeking greener solutions.

### Acknowledgements:

The authors thank the Department of Zoology at Ghazi University, Dera Ghazi Khan, Pakistan, for providing the necessary guidance.

### Ethical Considerations:

The study strictly adhered to national and institutional ethical standards for aquatic research. The Punjab Fisheries Department's Animal Ethics Committee approved all protocols (APF-188; 2023). All participating researchers provided verbal agreement before experimentation.

### Declarations:

#### Consent for Publication:

We, the authors of this review article, confirm that the work is original, has not been published elsewhere, and is not under consideration by another journal. All authors have reviewed and approved the final manuscript for submission.

#### Competing Interests:

The authors declare that they have no competing financial or other interests that could influence the work reported in this manuscript.

#### Funding:

The authors did not receive any funding.

#### Author Contributions:

A. H., and M. O. contributed to the work's concept and design. K. H. J. and A.S. were responsible for writing. M.O., H.A.A., N.J.H., and K.H.J., drafted the manuscript. All authors reviewed and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

## REFERENCES

Abdel-Ghany, H. M., El-Sisy, D. M., & Salem, M. E. S. (2023). A comparative study of effects of curcumin and its nanoparticles on the growth, immunity and heat stress resistance of Nile tilapia (*Oreochromis niloticus*). *Scientific Reports*, 13(1), 2523. <https://doi.org/10.1038/s41598-023-29343-z>

Abdelkarim, E. A., Elsamahy, T., El Bayomi, R. M., Hussein, M. A., Darwish, I. A., El-tahlawy, A. S., ... & Sobhy, M. (2025). Nanoparticle-driven aquaculture: transforming disease management and boosting sustainable fish farming practices. *Aquaculture International*, 33(4), 1-49. <https://doi.org/10.1007/s10499-025-01952-7>

Asad, F., Al Sulivany, B. S., Hassan, H. U., Nadeem, A., Rohani, M. F., Owais, M., ... & Arai, T. (2025). Evaluating the differential effect of growth and health parameters on *Oreochromis niloticus* and *Cirrhinus mrigala* under different rice protein concentration. *Egyptian Journal of Aquatic Research*, 51(1), 107-116. <https://doi.org/10.1016/j.ejar.2024.12.002>

Abdulrahman, P., & Al Sulivany, B. (2025). Dietary Quercus infectoria Mitigates Lead Nitrate Toxicity in Common Carp (*Cyprinus carpio*): Impacts on Growth Performance, Condition Factors, Weight Length Relationship, Hematological Responses, and Detoxification Potential During 60-Day Exposure. *Egyptian Journal of Aquatic Biology and Fisheries*, 29(2), 383-405. <https://doi.org/10.21608/ejabf.2025.416695>

Aebi, H. (1984). Catalase in vitro. In Methods in enzymology (Vol. 105, pp. 121-126). Academic press.

Al Sulivany, B. S., Gali Romani, F. A., Mohammed, D. A., & Khaleefah, R. S. (2024a). Winter Dietary Protein Impacts on Growth performance of *Cyprinus carpio*. *Egyptian Journal of Aquatic Biology & Fisheries*, 28(5). <https://doi.org/10.21608/ejabf.2024.380406>

Al Sulivany, B. S., Hassan, N. E., & Mhammad, H. A. (2024b). Influence of dietary protein content on growth performance, feed efficiency, condition factor, and length-weight relationship in *Cyprinus carpio* during the summer season. *Egyptian Journal of Aquatic Biology & Fisheries*, 28(2), 505-521. <https://doi.org/10.21608/ejabf.2024.349722>

Alsubaie, N., Abd-Elhakim, Y. M., Mohamed, A. A. R., Ibrahim, R. E., Metwally, M. M., Khamis, T., ... & Farag, M. F. (2025). Effects of purslane on hematology, *Aeromonas hydrophila* resistance, and NF-κB-LC3 pathway in Nile tilapia under waterborne cadmium. *Aquaculture International*, 33(6), 1-30. <https://doi.org/10.1007/s10499-025-02083-9>

Antache, A., Simionov, I. A., Petrea, S. M., Nica, A., Georgescu, P. L., Oprică, L., ... & Poroch, V. (2025). Insect-Antioxidants Symbiotic Nexus—Pathway for Sustainable and Resilient Aquaculture: A Case Study for Evaluating Koi Carp Growth and Oxidative Stress Status. *Antioxidants*, 14(4), 371. <https://doi.org/10.3390/antiox14040371>

Bhatt, D., Holeyappa, S. A., Pandey, A., Bansal, N., Hundal, J. S., & Khairnar, S. O. (2024). Growth performance, physiological response, and tissue microarchitecture of the carp *Labeo rohita* challenged with AFB1 are improved by supplementing with turmeric. *Spanish Journal of Agricultural Research*, 22(2), e0501-e0501. <https://doi.org/10.5424/sjar/202422-19974>

Bhattacharya, S., Das, B. K., Roy, A., Nayak, P. K., Saha, A., Parida, P. K., ... & Chakraborty, S. (2024). Socio-Economic Profile and Vulnerabilities Assessment in Small-Scale Fisheries of Lower Gangetic Flood Plain Wetland: An Initiative Towards Achieving Wetland Ecosystem Sustainability and Community Well-Being.

Sustainability, 16(21), 9583. <https://doi.org/10.3390/su16219583>

Bin-Ammar, A., Ahmeda, A. F., Abdelkarim, M., El-Bab, A. F. F., Amer, A. A., Abdelnour, S. A., ... & Naiel, M. A. (2024). Dietary curcumin nanoparticles improve growth performance, oxiDative status and immune response of european seabass (*Dicentrarchus labrax*). Ann. Anim. Sci., 24(3), 889-900. <https://doi.org/10.2478/aoas-2024-0018>

Bouyahya, A., Bakrim, S., Aboulaghra, S., El Kadri, K., Aanniz, T., Khalid, A., ... & El Omari, N. (2024). Bioactive compounds from nature: Antioxidants targeting cellular transformation in response to epigenetic perturbations induced by oxidative stress. Biomedicine & Pharmacotherapy, 174, 116432. <https://doi.org/10.1016/j.biopha.2024.116432>

Debnath, S., Karnataka, G., Puthiyottil, M., Lianthuamluuaia, L., Kumari, S., Das Ghosh, B., ... & Das, B. K. (2025). Improving approaches towards climate-smart wetland fisheries with a focus on small indigenous fishes. Sustainable Water Resources Management, 11(2), 1-11. <https://doi.org/10.1007/s40899-025-01198-4>

Díaz-Puertas, R., Rodríguez-Cañas, E., Lozoya-Agulló, M. J., Badía-Hernández, P. V., Álvarez-Martínez, F. J., Falcó, A., & Mallavia, R. (2024). Bovine serum albumin and lysozyme nanofibers as versatile platforms for preserving loaded bioactive compounds. International Journal of Biological Macromolecules, 280, 136019. <https://doi.org/10.1016/j.ijbiomac.2024.136019>

Eissa, E. S. H., Khattab, M. S., Elbahnaawy, S., Elshopakey, G. E., Alamoudi, M. O., Aljäbari, R. M., ... & Naiel, M. A. (2024). The effects of dietary *Spirulina platensis* or curcumin nanoparticles on performance, body chemical composition, blood biochemical, digestive enzyme, antioxidant and immune activities of *Oreochromis niloticus* fingerlings. BMC Veterinary Research, 20(1), 215. <https://doi.org/10.1186/s12917-024-04058-z>

El-Gogary, M. R., Dorra, T. M., & El-Sayed, I. A. (2025). Impact of dietary supplementation levels of turmeric powder (*curcuma longa*) on performance, carcass characteristics, blood biochemical, jejunum histological and gut microflora in broiler chickens. Journal of Animal and Poultry Production, 16(1), 7-13. <https://doi.org/10.21608/jappmu.2025.339434.1138>

Elkhaiat, I. A., El-Kassas, S., Abdo, S. E., El-Naggar, K., Shalaby, H. K., Nofal, R. Y., ... & Lestingi, A. (2025). Leverage of lysozyme dietary supplementation on gut health, hematological, antioxidant, and immune parameters in different plumage-colors Japanese quails. Poultry Science, 104(1), 104474. <https://doi.org/10.1016/j.psj.2025.104474>

Esmaelzadeh, N., Miri, M. S., Mavaddat, H., Peyrovinasab, A., Ghasemi Zargar, S., Sorous Kabiri, S., ... & Abdolghaffari, A. H. (2024). The regulating effect of curcumin on NF- $\kappa$ B pathway in neurodegenerative diseases: a review of the underlying mechanisms. Inflammopharmacology, 32(4), 2125-2151. <https://doi.org/10.1007/s10787-024-01482-3>

Fan, Z., Wang, Y., Li, C., Wu, D., Wang, L., Li, J., ... & Xu, H. (2025). Effects of acute cadmium exposure on biochemistry, liver oxidative stress and transcriptomics in Songpu mirror carp (*Cyprinus carpio* Songpu). Aquaculture Reports, 42, 102757. <https://doi.org/10.1016/j.aqrep.2025.102757>

Hafeez, A., Shahid Ali, S., Akhtar, J., Naz, S., Alrefaei, A. F., Albeshr, M. F., ... & Ullah Khan, R. (2024). Impact of coriander (*Coriandrum sativum*), garlic (*Allium sativum*), fenugreek (*Trigonella foenum-graecum*) on zootechnical performance, carcass quality, blood metabolites and nutrient digestibility in broilers chickens. Veterinary Quarterly, 44(1), 1-7. <https://doi.org/10.1080/01652176.2024.1234567>

Hematyar, N., Rahimnejad, S., Gorakh Waghmare, S., Malinovskyi, O., & Policar, T. (2024). Effects of stocking density and pre-slaughter handling on the fillet quality of largemouth bass (*Micropterus salmoides*): Implications for fish welfare. Foods, 13(10), 1477. <https://doi.org/10.3390/foods13101477>

Hoseini, S. M., & Al Sulivany, B. S. (2024). Copper And Microplastic Exposure Affects the Gill Gene Expression of Common Carp During Saltwater Challenge. Science Journal of University of Zakho, 12(3), 382-387. <https://doi.org/10.25271/sjuz.2024.12.3.1335>

Hossain, A., Habibullah-Al-Mamun, M., Nagano, I., Masunaga, S., Kitazawa, D., & Matsuda, H. (2022). Antibiotics, antibiotic-resistant bacteria, and resistance genes in aquaculture: risks, current concern, and future thinking. Environmental Science and Pollution Research, 1-22. <https://doi.org/10.1007/s11356-022-19050-z>

Jasim, S. A., Al-Dhalimy, A. M. B., Zokaei, M., Salimi, S., Alnajar, M. J., Kumar, A., ... & Gholami-Ahangaran, M. (2024). The Beneficial Application of Turmeric (*Curcuma longa L.*) on Health and Egg Production, in Layers: A Review. Veterinary Medicine and Science, 10(6), e70115. <https://doi.org/10.1002/vms3.70115>

Jomova, K., Alomar, S. Y., Alwasel, S. H., Nepovimova, E., Kuca, K., & Valko, M. (2024). Several lines of antioxidant defense against oxidative stress: Antioxidant enzymes, nanomaterials with multiple enzyme-mimicking activities, and low-molecular-weight antioxidants. Archives of Toxicology, 98(5), 1323-1367. <https://doi.org/10.1007/s00204-024-03696-4>

Kasihmuddin, S. M., Cob, Z. C., Noor, N. M., & Das, S. K. (2024). Effect of different temperature variations on the physiological state of catfish species: A systematic review. Fish Physiology and Biochemistry, 50(2), 413-434. <https://doi.org/10.1007/s10695-024-01319-4>

Khaldoun, H., Settar, A., Oularbi, Y., Boudjema, N., Amokrane, A., Djennane, N., & Tarzaali, D. (2025). The effect of thyme essential oil on duodenal toxicity induced by subacute exposure to Voliam Targo insecticide in male rabbits. Toxicology Reports, 14, 101959. <https://doi.org/10.1016/j.toxrep.2025.101959>

Khanjani, M. H., Sharifinia, M., & Emerenciano, M. G. C. (2024). Biofloc technology (BFT) in aquaculture: What goes right, what goes wrong? A scientific-based snapshot. Aquaculture Nutrition, 2024(1), 7496572. <https://doi.org/10.1155/2024/7496572>

Khieokhajonkhet, A., Roatboonsongsri, T., Suwannalers, P., Aeksiri, N., Kaneko, G., Ratanasut, K., ... & Phromkunthong, W. (2023). Effects of dietary supplementation of turmeric (*Curcuma longa*) extract on

growth, feed and nutrient utilization, coloration, hematology, and expression of genes related immune response in goldfish (*Carassius auratus*). *Aquaculture Reports*, 32, 101705. <https://doi.org/10.1016/j.aqrep.2023.101705>

Komal, W., Fatima, S., Minahal, Q., & Liaqat, R. (2024). Enhancing growth, antioxidant capacity, and immune response in tilapia (*Oreochromis niloticus*) through curcumin supplementation across varied stocking density paradigms. *PLoS ONE*, 19(11), e0311146. <https://doi.org/10.1371/journal.pone.0311146>

Kusi, I. O., Obirikorang, K. A., & Adjei-Boateng, D. (2025). Effects of phytobiotic *Curcuma longa*, *Allium sativum* and *Zingiber officinale*-supplemented diets on growth, utilisation of feed and Nile tilapia (*Oreochromis niloticus*) resistance against *Streptococcus agalactiae*. *Aquaculture Research*, 2025(1), 9911375. <https://doi.org/10.1155/2025/9911375>

Lulijwa, R., Alfaro, A. C., Merien, F., Burdass, M., Venter, L., & Young, T. (2019). In vitro immune response of chinook salmon (*Oncorhynchus tshawytscha*) peripheral blood mononuclear cells stimulated by bacterial lipopolysaccharide. *Fish & shellfish immunology*, 94, 190-198. DOI: 10.1016/j.fsi.2019.07.063

Marklund, S., & Marklund, G. (1974). Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry*, 47(3), 469-474. <https://doi.org/10.1111/j.1432-1033.1974.tb03714.x>

Meitei, M. M., Patel, A. B., Singh, S. K., Biswas, P., Behera, S. K., & Upadhyay, A. D. (2025). Is individual or combination of three Indian major carps (Catla catla, *Labeo rohita*, and *Cirrhinus mrigala*) preferable secondary co-species in semi-intensive polyculture with pabda (*Ompok bimaculatus*)? *Aquaculture International*, 33(6), 1-23. <https://doi.org/10.1007/s10499-025-01234-2>

Metwaly, S., Nasr, H., Ahmed, K., & Fathi, M. (2025). Multifaceted stress response in Nile tilapia (*Oreochromis niloticus*) fingerlings: Integrative analysis of salinity, ammonia, and stocking density effects on growth, physiology, and gene expression. *Fish Physiology and Biochemistry*, 51(1), 1-10. <https://doi.org/10.1007/s10695-024-01319-4>

Mithuna, R., Tharanyalakshmi, R., Jain, I., Singhal, S., Sikarwar, D., Das, S., ... & Das, B. (2024). Emergence of antibiotic resistance due to the excessive use of antibiotics in medicines and feed additives: A global scenario with emphasis on the Indian perspective. *Emerging Contaminants*, 10, 100389. <https://doi.org/10.1016/j.emcon.2024.100389>

Munguti, J., Muthoka, M., Chepkirui, M., Kyule, D., Obiero, K., Ogello, E., ... & Kwikiriza, G. (2024). The fish feed sector in Kenya, Uganda, Tanzania, and Rwanda: Current status, challenges, and strategies for improvement—A comprehensive review. *Aquaculture Nutrition*, 2024(1), 8484451. <https://doi.org/10.1155/2024/8484451>

Nabi, N., Ahmed, I., Qadir, M., & Reshi, Q. M. (2025). Global aquaculture: Scenarios and nutritional implications. In *Aquaculture: Enhancing food security and nutrition* (pp. 121-140). Springer. [https://doi.org/10.1007/978-3-031-55886-9\\_6](https://doi.org/10.1007/978-3-031-55886-9_6)

Naghdi, S., Rezaei, M., Heidari, M. G., Tahergorabi, R., Lorenzo, J. M., & Mirzaei, F. (2024). Insights into fishery by-product application in aquatic feed and food: A review. *Aquaculture International*, 32(5), 5851-5910. <https://doi.org/10.1007/s10499-024-01476-6>

Neil, B., Cheney, G. L., Rosenzweig, J. A., Sha, J., & Chopra, A. K. (2024). Antimicrobial resistance in aeromonads and new therapies targeting quorum sensing. *Applied Microbiology and Biotechnology*, 108(1), 205. <https://doi.org/10.1007/s00253-024-13080-y>

Owais, M., Al Sulivany, B., Abdulhalim, B. A., & Mehroz, R. (2024). The Pangas catfish *Pangasius pangasius*: Growth efficiency and nutritional composition under variety of saltwater challenges. *Egyptian Journal of Aquatic Biology & Fisheries*, 28(6), 123-140. <https://doi.org/10.21608/ejabf.2024.354678>

Owais, M., Fazal, R. M., Riaz-Ud-Din, Q. U. R. E. S. H. I., Yasin, R., Irfan, M., Ameer, M. W., & Sarwar, A. (2023). Evaluation of different salinity levels on growth performance and proximate composition of *Wallago attu*. *University of Sindh Journal of Animal Sciences (USJAS)*, 7(2), 44-49. <https://doi.org/10.46610/USJAS.2023.v07i02.004>

Pitiot, A., Rolin, C., Seguin-Devaux, C., & Zimmer, J. (2025). Fighting antibiotic resistance: Insights into human barriers and new opportunities. *BioEssays*, \*46\*(3), e70001. <https://doi.org/10.1002/bies.202370001>

Rasheed, H. A., Rehman, A., Karim, A., Al-Asmari, F., Cui, H., & Lin, L. (2024). A comprehensive insight into plant-derived extracts/bioactives: Exploring their antimicrobial mechanisms and potential for high-performance food applications. *Food Bioscience*, \*58\*, 104035. <https://doi.org/10.1016/j.fbio.2024.104035>

Rasul, M. N., Hossain, M. T., Haider, M. N., Hossain, M. T., & Reza, M. S. (2025). Disease prevalence, usage of aquaculture medicinal products and their sustainable alternatives in freshwater aquaculture of North-Central Bangladesh. *Veterinary Medicine and Science*, \*11\*(2), e70276. <https://doi.org/10.1002/vms3.70276>

Reshi, Q. M., Ahmad, I., Nabi, N., Ahmed, I., & Sidiq, M. (2025). Global food insecurity: Role of fishes to compensate food insecurity and nutritional demand. In *Food security, nutrition and sustainability through aquaculture technologies* (pp. 73-93). Springer Nature. [https://doi.org/10.1007/978-3-031-55886-9\\_5](https://doi.org/10.1007/978-3-031-55886-9_5)

Roney, M., Huq, A. M., Rullah, K., Zamri, N. B., & Mohd Aluwi, M. F. F. (2025). Curcumin, a bioactive compound of turmeric (*Curcuma longa*) and its derivatives as  $\alpha$ -amylase and  $\alpha$ -glucosidase inhibitors. *Cell Biochemistry and Biophysics*, \*83\*(1), 53-71. <https://doi.org/10.1007/s12013-024-01337-2>

Shin, H. Y., Kim, Y. S., Shin, K. S., Han, S. H., Suh, H. J., & Yu, K. W. (2024). Effect of rhamnogalacturonan-I-rich polysaccharides isolated from crabapple hydrolysates on IL-1 $\beta$ -induced inflammation in intestinal epithelial cells. *International Journal of Biological Macromolecules*, \*277\*, 134240. <https://doi.org/10.1016/j.ijbiomac.2024.134240>

Song, X., Kong, F., Liu, B. F., Song, Q., Ren, N. Q., & Ren, H. Y. (2024). Antioxidants alleviated low-temperature stress in microalgae by modulating reactive oxygen species to improve lipid production and antioxidant defense. *Bioresource Technology*, \*413\*, 131451. <https://doi.org/10.1016/j.biortech.2024.131451>

Tahir, R., Afzal, F., Ghaffar, A., Liang, J., Shrestha, A., Habiba, U., & Yang, S. (2024). Effectiveness of lycopene dietary enrichment for growth performance, digestive enzymes, blood health, immunity, and antioxidant status of common carp (*Cyprinus carpio*) against chronic glyphosate toxicity. *Aquaculture*, \*590\*, 741084. <https://doi.org/10.1016/j.aquaculture.2024.741084>

Taysi, M. R., Kirici, M., Kirici, M., Tuzun, B., & Poustforoosh, A. (2024). Antioxidant enzyme activities, molecular docking studies, MM-GBSA, and molecular dynamic of chlorpyrifos in freshwater fish *Capoeta umbla*. *Journal of biomolecular structure and dynamics*, 42(1), 163-176. [DOI: 10.1080/07391102.2023.2196698](https://doi.org/10.1080/07391102.2023.2196698)

Thavorn, K., Wolfe, D., Faust, L., Shorr, R., Akkawi, M., Isaranuwatchai, W., ... & Hutton, B. (2024). A systematic review of the efficacy and safety of turmeric in the treatment of digestive disorders. *Phytotherapy Research*, 38(6), 2687-2706. <https://doi.org/10.1002/ptr.8189>

Verdi, C. M., Baldissera, M. D., Machado, V. S., Milarch, C. F., de Andrade, E. N. C., Dantas, G. A., ... & Santos, R. C. V. (2024). Gaultheria procumbens essential oil longevity and oxidant/antioxidant status of silver catfish *Rhamdia quelen* experimentally infected by *Aeromonas caviae*. *The Microbe*, 5, 100191. [DOI: 10.1016/j.microb.2024.100191](https://doi.org/10.1016/j.microb.2024.100191)

Wang, Y., Wang, L., Zhang, H., Ren, P., Cheng, X., Hong, F., ... & Gou, D. (2025). Immunostimulatory effects mechanism of polysaccharide extracted from *Acanthopanax senticosus* on RAW 264.7 cells through activating the TLR/MAPK/NF-κB signaling pathway. *Scientific Reports*, 15(1), 13440. <https://doi.org/10.1038/s41598-025-97423-3>

Wei, L. S., Hooi, K. Y., Khoo, M. I., Mn, A., & Wee, W. (2024). Effect of dietary supplementation of turmeric, *Curcuma longa* leaf on growth and health status of African catfish, *Clarias gariepinus*: *Veterinary Integrative Sciences*, 22(3), 907-920. <https://doi.org/10.12982/VIS.2024.062>

Wildhaber, M. L., West, B. M., Ditter, K. K., Moore, A. P., & Peterson, A. S. (2023). A review of grass carp and related species literature on diet, behavior, toxicology, and physiology focused on informing development of controls for invasive grass carp populations in North America. *Fishes*, 8(11), 547. <https://doi.org/10.3390/fishes8110547>

Yadav, N. K., Patel, A. B., Singh, S. K., Mehta, N. K., Anand, V., Lal, J., ... & Devi, N. C. (2024). Climate change effects on aquaculture production and its sustainable management through climate-resilient adaptation strategies: A review. *Environmental Science and Pollution Research*, 31(22), 31731-31751. <https://doi.org/10.1007/s11356-024-33435-2>

Zheng, P., Zhou, B., Zheng, Y., Ke, Y., Zhang, W., Xing, M., & Zhao, H. (2025). Unveiling the toxic mechanism of arsenic on the spleen of *Cyprinus carpio* and the antagonistic role of zinc via the P38 MAPK/Nrf2/HO-1 pathway. *Biological Trace Element Research*. Advance online publication. <https://doi.org/10.1007/s12011-025-04045-x>

Zou, C., Jiang, H., Wu, X., Gao, J., & Ma, W. (2025). Integrative analysis of metabolomics and transcriptomics reveals alterations in egg quality and hepatic lipid metabolism in hens supplemented with curcumin. *Animal Nutrition*, 21, 302-315. <https://doi.org/10.1016/j.aninu.2025.01.005>