

Original article

Growth, Antioxidant Defense, and Immune Response in *Labeo rohita* Fingerlings with the Dietary Vitamin C Supplementation

Zainab Asghar¹, Nuha Hameed Albassam², Milat Hussain Mustafa³, Hifza Afzaal⁴, Zeliha Selamoglu^{5,6,7*}, Arooj Fatima⁸, Danish kamal⁹, Rashid Hussain¹⁰,

¹Department of Zoology, Lahore College for Women University, Lahore, Punjab, Pakistan.

²Department of Animal Production, College of Agriculture, University of Tikrit, Salah Al-deen, Iraq

³Department of Radiological Imaging Technology, Cihan University-Duhok, Duhok, Kurdistan Region, Iraq

⁴Department of Zoology, University of Education, Township, Lahore, Punjab, Pakistan.

⁵Department of Medical Biology, Medicine Faculty, Nigde Omer Halisdemir University, Nigde, Türkiye.

⁶Western Caspian University, Baku, Azerbaijan.

⁷Department of Biology, Faculty of Sciences, Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan.

⁸Department of Zoology, Government College University Lahore, Punjab, Pakistan.

⁹State Key Laboratory of Tea Plant Germplasm Innovation and Resource Utilization, Anhui Agricultural University, Hefei 230036, Anhui, China.

¹⁰Department of Zoology, Ghazi University, Dera Ghazi Khan, Punjab, Pakistan.

*Corresponding author: zselamoglu@ohu.edu.tr

ABSTRACT

Received:
24, Dec, 2025

The study was designed to investigate the vitamin C supplementation effect on growth, oxidative stress, and immune response of *Labeo rohita*. Four diets 0, 200, 400, and 800 mg/kg was designed, 480 fingerlings with the initial weight 6.11 ± 0.08 g in triplicate groups were randomly distributed into 12 fiberglass tanks for 60 days. Growth results showed that final weight (13.87 ± 0.23 – 19.83 ± 0.34 g), weight gain (152.6 ± 2.7 – $251.1 \pm 3.5\%$), and specific growth rate (1.91 ± 0.06 – $2.52 \pm 0.03\%$ /day) increased significantly ($p < 0.05$) with increasing dietary vitamin C levels, while feed conversion ratio (1.81 ± 0.04 – 1.29 ± 0.04) decreased ($p = 0.006$), indicating improved feed utilization.

Accepted:
17, Jan, 2026

Survival rates were $91.1 \pm 2.4\%$ in the control, whereas they reached $98.4 \pm 1.4\%$ in the 800 mg/kg group. Antioxidant enzyme showed a significant ($p < 0.01$) enhancement. Superoxide dismutase (SOD) and catalase (CAT) activities increased from 25.6 ± 1.3 to 36.1 ± 1.3 U/mg protein and 12.4 ± 0.7 to 20.1 ± 0.5 U/mg protein. The malondialdehyde (MDA) levels reduced significantly ($p = 0.005$), while glutathione (GSH) concentration increased significantly ($p = 0.007$) from 3.10 ± 0.13 to 4.41 ± 0.13 $\mu\text{mol/g}$ tissue. Immune response parameters were also positively affected by dietary vitamin C. Lysozyme activity increased significantly ($p = 0.008$) from 15.4 ± 0.3 to 23.4 ± 0.6 U/mL, respiratory burst activity improved from 0.178 ± 0.006 to 0.261 ± 0.006 OD₆₃₀ nm, and phagocytic activity rose from 28.3 ± 1.4 to $41.4 \pm 1.4\%$. Similarly, nitric oxide production increased from 4.13 ± 0.28 to 6.86 ± 0.26 $\mu\text{mol/L}$. Total protein, albumin, and globulin were also significantly raised ($p < 0.05$). The study concluded that dietary vitamin C can be used to increase growth, support antioxidant defense, and improve immune responses in *L. rohita*.

KEYWORDS: *Labeo rohita*, Vitamin C, Growth performance, Oxidative stress, Immune response.

1. INTRODUCTION

Aquaculture in many developing regions fastest-expanding sector, contributing significantly to global food production,

economic development, nutritional security, and livelihood support (Irshath *et al.*, 2023). In South Asia, aquaculture, predominantly in Pakistan, India, and Bangladesh, relies heavily on inland fish production, including *Labeo rohita*, *Catla catla*, and *Cirrhinus mrigala* (Abdul-Qadir *et al.*, 2024). Among these,

Access this article online



<https://doi.org/10.25271/sjuz.2026.14.1.1860>

Printed ISSN 2663-628X;
Electronic ISSN 2663-6298

Science Journal of University of Zakho
Vol. 14, No. 01, pp. 164–170 January-2026

This is an open access under a CC BY-NC-SA 4.0 license
(<https://creativecommons.org/licenses/by-nc-sa/4.0/>)

L. rohita is recognized as the most commercially valuable freshwater species, with adaptability to semi-intensive and intensive culture systems and a fast growth rate (Gupta *et al.*, 2021). However, high stocking densities increased feeding rates, feed efficiency, and survival in aquaculture, but imperfect water exchange disrupted immunity, caused physiological stress, and increased susceptibility to infectious disease (Hossain *et al.*, 2022; Namiq *et al.*, 2025). One of the most effective ways to reduce such adverse effects is through nutritional changes, such as integrating micronutrients and antioxidants into the diet, including vitamins and minerals that play critical roles in maintaining metabolic function, immune function, and oxidative balance (Ibraheem *et al.*, 2017; Awuchi *et al.*, 2022). Among these, vitamin C (ascorbic acid), due to its multifaceted physiological functions responsible for synthesizing glucose in the liver, maintaining health and performance in cultured fish species, and interpreting them as completely dependent on dietary sources of vitamin C (Jiménez *et al.*, 2018). Vitamin C directly interacts synergistically with lipid-soluble antioxidants and reactive oxygen species with free radicals, allowing for the protection of cellular membranes from oxidative damage (Soldado *et al.*, 2021). Furthermore, vitamin C increases the activities of antioxidant enzymes, such as SOD, catalase, and glutathione peroxidase, and minimizes oxidative injury in tissues (El-Gendy *et al.*, 2010). The disparity between the making of ROS and the antioxidant defense system leads to oxidative stress, subsequent protein denaturation, and lipid peroxidation that impairs growth and biological performance; however, the vitamin C supplementation in fish diets progresses antioxidant enzyme activities, improving oxidative stress in *Oreochromis niloticus* (Surai *et al.*, 2019) and *Clarias gariepinus* (Abalaka *et al.*, 2021).

The antioxidative role of vitamin C is closely linked to the enhancement of immune responses against invading pathogens in fish, and its efficacy is also influenced by nutritional status (Ibrahim *et al.*, 2021). Vitamin C supplementation modulates immune parameters, including lysozyme activity, respiratory burst, nitric oxide production, and serum protein profiles (Firouzi *et al.*, 2022). Furthermore, it also enhances macrophage and neutrophil activity and enhances antibody production (Zhang *et al.*, 2023). The dietary intake of vitamin C enhances resistance to bacterial infections, increasing lysozyme and respiratory burst activities in freshwater fish (Xue *et al.*, 2022). Moreover, vitamin C plays an essential role in reducing immunosuppression induced by ecological or nutritional stressors by maintaining the functional integrity of the immune system, and it also interacts with other dietary components that help regulate cellular signaling pathways involved in immune parameters (Tang *et al.*, 2024). These nutritional formulations with vitamin C are important, especially at the fingerling stage, for the rapid growth, metabolic activity, and greatly enhance survival under intensive culture systems (Xu *et al.*, 2022). Despite significant progress in nutritional physiology, the rarity persists in dietary vitamin C data on growth, oxidative stress biomarkers, and immune responses of *Labeo rohita* fingerlings. Thus, the present study was directed to assess the outcome of dietary vitamin C

supplementation on growth, oxidative stress biomarkers, and immune response parameters in *L. rohita* fingerlings.

2. MATERIALS AND METHODS

Experimental Design:

This present study was planned to examine the outcome of dietary vitamin C on the growth, oxidative stress biomarkers, and immune response of *L. rohita* fingerlings. 480 total of *L. rohita* fingerlings with an average initial weight of 6.112 ± 0.10 g were purchased from the Government Hatchery, Lahore, Punjab, Pakistan. Laboratory conditions for a period of 14 days were provided to the experimental fish, which were fed a basal diet twice daily during this period, and acclimated in fiberglass tanks (400 L capacity of water). The fingerlings, after acclimation, were randomly distributed among 12 fiberglass tanks at a stocking density of 40 fish per tank. Four dietary treatments with triplicate were designed, viz, control diet (0 mg/kg vitamin C) and three diets supplemented with 200, 400, and 800 mg/kg vitamin C, respectively. Siphoning (20% of the water removed from each tank) was performed daily to remove debris and uneaten feed throughout the experiment.

The fish were fed at 5% of their body weight for 60 days, the water parameters were recorded such as temperature; $26.5 \pm 0.4^\circ\text{C}$, dissolved oxygen; 7.12 ± 0.06 mg/L, and pH; 7.98 ± 0.02 and total ammonia nitrogen (TAN) was maintained through regular water exchange below 0.05 mg/L, while nitrite (NO_2^-) levels remained below 0.1 mg/L throughout the experimental period. Continuous aeration was provided using air stones connected to a central air compressor. After the end of the study period, the experimental fish were fasted for 24 hours before sampling. 40 fish from each group were randomly selected for both the biochemical and growth analyses. The clove oil (50 mg/L) was used, and fish were anesthetized, dissection performed, liver tissue was removed and controlled in ice-cold phosphate buffer, and centrifuged at 4°C . Blood samples were collected using sterile heparinized syringes from the caudal vein, centrifuged at 3,000 rpm for 10 min at 4°C to obtain serum, and stored at -20°C until analysis of immune response parameters. The protocol was conducted in accordance with subsequent ethical standards for fish nutrition research (Mohamad Hassan *et al.*, 2025).

Dietary Preparation:

The study used the following ingredients: fish meal, soybean meal, maize gluten, rice bran, and wheat flour, which were finely crushed and carefully mixed using an automatic mixer. Vitamin C (It was provided in the diet in the form of ascorbyl-2-polyphosphate) at inclusion levels of 0, 200, 400, and 800 mg/kg diet was combined into the basal diet. After the dry mixing, fish oil and tolerable purified water were added to get a consistent dough appropriate for pelleting. The laboratory pellet machine equipped with a 2-mm die was used to produce uniform pellets. The pellets were air-dried at room temperature ($28\text{--}30^\circ\text{C}$) for 48 hours, followed by storage in airtight containers at 4°C until further use to preserve nutrient veracity. The comprehensive composition of experimental diets is presented in Table 1.

Table 1: The diet composition for *Labeo rohita* fingerlings

Ingredients (g/kg)	Control (0 mg/kg Vit C)	Vit C (200 mg/kg)	Vit C (400 mg/kg)	Vit C (800 mg/kg)
Fish meal	200	200	200	200
Soybean meal	300	300	300	300
Maize gluten meal	120	120	120	120
Rice bran	150	150	150	150
Wheat flour	150	150	150	150
Fish oil	50	50	50	50
Vitamin–mineral premix¹	20	20	20	20
Dicalcium phosphate	5	5	5	5
Vitamin C (mg/kg)²	–	200	400	800
Total	1000	1000	1000	1000

Note: Vitamin–mineral premix such as vitamin A, vitamin D₃, vitamin E, vitamin K₃, vitamin B₁, vitamin B₂, vitamin B₆, vitamin B₁₂, niacin, folic acid, biotin, calcium pantothenate, choline chloride, Fe, Zn, Mn, Cu, I, and Se.

Growth Performance Metrics:

Growth parameters evaluation performed by using the following standard formulas:

$$WG, \% = [(FBW - IBW) / IBW] \times 100$$

$$SGR, \%/\text{day} = [(\ln FBW - \ln IBW) / \text{Duration (days)}] \times 100$$

$$FCR = FI (\text{g}) / WG (\text{g})$$

$$\text{Survival rate (\%)} = (\text{Final fish number} / \text{Initial fish number}) \times 100.$$

Antioxidant Enzyme Activities:

The antioxidant enzymatic markers were examined, such as the Superoxide dismutase (SOD) activity recorded by the inhibition of pyrogallol auto-oxidation at 560 nm spectrophotometrically. Catalase (CAT) activity was examined by the rate of hydrogen peroxide decomposition with recorded absorbance at 240 nm. Lipid peroxidation was estimated by malondialdehyde (MDA) levels by (TBARS) assay, at 532 nm. GSH (Reduced Glutathione) concentration was determined according to Ilhan *et al.* (2004). Total Protein content was assessed using the method of Moore *et al.* (2010).

Immune Response Assays:

Immune parameters were analyzed using serum samples obtained from blood centrifugation and stored at -20 °C until analysis. Lysozyme activity was determined using the turbidimetric method described by Fazal *et al.* (2025) with *Micrococcus lysodeikticus* as the substrate, and results were expressed as U/mL serum based on the reduction in absorbance

at 530 nm. Respiratory burst activity was measured following Rothe *et al.* (1991) using the nitroblue tetrazolium (NBT) reduction assay and expressed as optical density at 600 nm. Phagocytic activity was assessed according to Pavlou *et al.* (2017) and expressed as percentage phagocytosis. Nitric oxide (NO) production was measured in serum using the Griess reagent, following Privat *et al.* (1997), with absorbance read at 540 nm and results expressed as μmol/mL. Serum total protein, albumin, and globulin concentrations were determined using the Biuret method, bromocresol green method, and calculated difference, respectively, following Katsoulos *et al.* (2017), and expressed as g/dL.

3. RESULTS

The vitamin C had a significant effect on the growth performance of *L. rohita* (Table 2). The IW of fish did not fluctuate significantly ($p > 0.05$) among the treatments. The FW was 13.87 ± 0.23 g in the control, 19.83 ± 0.34 g in the 800 mg/kg vitamin C group, which was significantly increased ($p = 0.007$). Similarly, WG at the highest vitamin C level was $251.1 \pm 3.5\%$ and lowest in the control ($152.6 \pm 2.7\%$) ($p = 0.008$). The SGR was $1.91 \pm 0.06\%/\text{day}$ in the control group, which improved to $2.52 \pm 0.03\%/\text{day}$ in the 800 mg/kg group significantly ($p = 0.007$). The FCR in the control group was 1.81 ± 0.04 , whereas a gradual decline was observed in the 400 and 800 mg/kg vitamin C diets, with FCR values of 1.34 ± 0.06 and 1.29 ± 0.04 , respectively ($p = 0.006$). The survival rate was $91.1 \pm 2.4\%$ in the control group and $98.4 \pm 1.4\%$ in the 800 mg/kg vitamin C group, which indicated significant ($p = 0.006$) improvement.

Table 2: The measurements of growth parameters on different grades of vitamin C supplementation in *Labeo rohita* fingerlings.

Parameters	Control (0 mg/kg Vit C)	Vit C (200 mg/kg)	Vit C (400 mg/kg)	Vit C (800 mg/kg)	p-value
IW (g)	6.11± 0.08	6.13± 0.06	6.14 ± 0.07	6.12± 0.05	NS
FW (g)	13.87 ± 0.23 ^d	15.73 ± 0.34 ^c	17.83± 0.42 ^b	19.83 ± 0.34 ^a	0.007
WG (%)	152.6± 2.7 ^d	189.7± 3.4 ^c	233.6± 4.6 ^b	251.1 ± 3.5 ^a	0.008
SGR (%/day)	1.91± 0.06 ^d	2.16± 0.05 ^c	2.44± 0.04 ^b	2.52± 0.03 ^a	0.007
FCR	1.81± 0.04 ^a	1.51± 0.06 ^b	1.34± 0.06 ^c	1.29± 0.04 ^c	0.006
SR (%)	91.1± 2.4 ^b	95.4± 1.7 ^a	96.4± 1.8 ^a	98.4 ± 1.4 ^a	0.006

The effect of vitamin C on the oxidative stress biomarkers of fingerlings is presented in Table 3. The SOD activity was 36.1 ± 1.3 U/mg protein in the 800 mg/kg vitamin C group, while 25.6 ± 1.3 U/mg protein was recorded in the control group (p = 0.006). Similarly, CAT activity was 20.1 ± 0.5 U/mg protein in the 800 mg/kg group, whereas 12.4 ± 0.7 U/mg protein was observed in the control group, which showed a significant (p = 0.007) enhancement. In contrast, the MDA concentration was 6.33 ±

0.24 nmol/mg protein in the control group, but 3.27 ± 0.17 nmol/mg protein was recorded in the group receiving 800 mg/kg vitamin C. This decrease in MDA values reflected a reduced oxidative damage and cellular lipid peroxidation. GSH levels increased significantly (p = 0.007): in the 800 mg/kg group, 4.41 ± 0.13 μmol/g tissue; and in the control group, 3.1 ± 0.13 μmol/g tissue.

Table 3: The measurement of oxidative stress biomarkers at various vitamin C levels in *L. rohita*.

Parameters	Control (0 mg/kg Vit C)	Vit C (200 mg/kg)	Vit C (400 mg/kg)	Vit C (800 mg/kg)	p-value
SOD, U/mg protein	25.6±1.3 ^c	31.6±1.2 ^b	34.7± 1.4 ^a	36.1±1.3 ^a	0.006
CAT, U/mg protein	12.4±0.7 ^c	16.3±0.6 ^b	17.6 ± 0.6 ^a	20.1±0.5 ^a	0.007
MDA, nmol/mg protein	6.33±0.24 ^a	4.21±0.12 ^b	3.44± 0.16 ^c	3.27±0.17 ^c	0.005
GSH, μmol/g tissue	3.1±0.13 ^c	3.56±0.12 ^b	4.34± 0.02 ^a	4.41±0.13 ^a	0.007

SOD; for Superoxide dismutase (U/mg protein), CAT; For Catalase (U/mg protein), MDA; for malondialdehyde (nmol/mg protein), GSH; Glutathione peroxidase (μmol/g tissue).

Dietary supplementation with vitamin C on *L. rohita* revealed a significant (p < 0.01) effect on non-specific immune responses. Lysozyme activity was 15.4 ± 0.3 U/mL in the control group and 23.4 ± 0.6 U/mL in the 800 mg/kg group. The respiratory burst activity was significantly enhanced compared with the control group, from 0.178 ± 0.006 to 0.261 ± 0.006 OD₆₃₀ nm at the highest vitamin C level. The Phagocytic activity was 28.3 ± 1.4% in the control group and 41.4 ± 1.4% in the 800

mg/kg group. Nitric oxide concentration rising from 4.13 ± 0.28 μmol/L in the control to 6.86 ± 0.26 μmol/L in the highest supplemented group (p = 0.005). Total protein improved from 3.15 ± 0.03 g/dL in the control up to 4.01 ± 0.05 g/dL, while albumin and globulin also enhanced significantly. The highest albumin level (1.59 ± 0.02 g/dL) and globulin level (2.41 ± 0.06 g/dL) were detected in the 800 mg/kg group.

Table 4: The evaluation of non-specific immune responses and serum biochemical parameters at various levels of vitamin in *L. rohita*.

Parameters	Control (0 mg/kg Vit C)	Vit C (200 mg/kg)	Vit C (400 mg/kg)	Vit C (800 mg/kg)	p-value
Lysozyme activity (U/mL)	15.4± 0.3 ^d	18.6± 0.7 ^c	22.4± 0.6 ^b	23.4± 0.6 ^a	0.008
Respiratory burst activity (OD₆₃₀ nm)	0.178 ± 0.006 ^d	0.211± 0.005 ^c	0.239± 0.007 ^b	0.261± 0.006 ^a	0.007
Phagocytic activity (%)	28.3± 1.4 ^d	31.9± 1.3 ^c	38.7± 1.6 ^b	41.4± 1.4 ^a	0.007
Nitric oxide (μmol/L)	4.13± 0.28 ^d	5.31± 0.21 ^c	6.41± 0.23 ^b	6.86± 0.26 ^a	0.005
Total protein (g/dL)	3.15 ± 0.03 ^d	3.31± 0.05 ^c	3.76± 0.06 ^b	4.01± 0.05 ^a	0.006
Albumin (g/dL)	1.31± 0.03 ^d	1.37± 0.06 ^c	1.55± 0.08 ^b	1.59± 0.02 ^a	0.006

Globulin (g/dL)	1.71± 0.04 ^d	2.23± 0.06 ^c	2.41 ± 0.05 ^b	2.41 ± 0.06 ^a	0.006
------------------------	-------------------------	-------------------------	--------------------------	--------------------------	-------

4. DISCUSSION

The improvement in growth parameters indicates a positive effect on growth and feed efficiency at higher vitamin C levels. Ibrahim *et al.* (2020) studied *Oreochromis niloticus* and described the effect of vitamin C on growth performance, collagen synthesis, and connective tissue development. Narra *et al.* (2015)

described that vitamin C in *Clarias gariepinus* increases the metabolic processes and growth parameters, with the muscle development and tissue repair. Fasil *et al.* (2021) reported that dietary vitamin C supplementation reduced oxidative stress and improved nutrient utilization, thereby improving growth outcomes. The present findings are also similar to Abdelwahab *et al.* (2020), who described that ascorbic acid controls endocrine and metabolic functions, promotes appetite, and protein combination in sea bass.

This study described the improved feed conversion ratio with vitamin C. Similarly, Magouz *et al.* (2020) also studied the enhanced digestive efficiency, improved FCR by stimulating enzyme activity that leads to better nutrient absorption and energy utilization. Increasing vitamin C supplementation increases survival by strengthening physiological tolerance and reducing stress-induced mortality. This observation agrees with Liua *et al.* (2016), who studied that vitamin C improves the protection of fish resilience against pathogens and stress, with the maintenance of the survival rate. Biswas *et al.* (2020) studied that vitamin C regenerates other antioxidants and neutralizes reactive oxygen species, and increasing the survival rate. In the present study, enhanced enzymatic activities with vitamin C thereby mitigating oxidative damage. Similar trends were reported by Daniel *et al.* (2021), who found that vitamin C supplementation significantly increased SOD and CAT activities in *Pangasius pangasius*, indicating oxidative stress. Moreover, elevated vitamin C concentrations led to a significant decrease in malondialdehyde levels, indicating reduced oxidative tissue damage and lipid peroxidation in *L. rohita*. This decrease in MDA is consistent with earlier findings in *Oreochromis niloticus* and *Cyprinus carpio*, in which a vitamin C supplement protects membrane lipids from degradation and reduces lipid peroxidation (Sherif *et al.*, 2024). The GSH, as a major non-enzymatic antioxidant, increased with the increase of vitamin C, which supports the maintenance of redox balance and detoxification of free radicals. These results agree with Singh *et al.* (2017), who reported that dietary vitamin C enhances GSH synthesis, thereby supporting the defense of cells against oxidative injury and antioxidant capacity. Vitamin C also improves both humoral and cellular immune mechanisms in *L. rohita* in the present study which is align with findings by Misra *et al.* (2007) who described the enhanced lysozyme and increased respiratory burst activities in fish fed vitamin-supplemented diets. Alberts *et al.* (2025) studied the higher macrophage, oxide production, and neutrophil activity with improved pathogen-killing ability and leukocyte function. Nitric oxide, with a higher level during immune responses, strengthened immune functionality (Gao *et al.*, 2021). The improvement in serum biochemical parameters with vitamin C regarded as important indicators of immune and nutritional

status. The total protein and globulin levels improved the synthesis of immunoglobulins and improved immune balance. Taalab *et al.* (2022) designated improved serum protein with the metabolic efficiency in *Clarias gariepinus* with vitamin C supplementation.

CONCLUSION

The present findings indicate that dietary vitamin C increases growth performance, decreases oxidative stress, and reinforces the immune system of *L. rohita* fingerlings, and suggest that vitamin C supplementation at 800 mg/kg is necessary to improve growth, health, and immune competence in *L. rohita*.

Acknowledgments:

The authors wish to express their sincere gratitude to the Department of Zoology, Lahore College for Women University, for their invaluable institutional support.

Ethical Statement:

The study protocol was approved by the, Lahore College for Women University (Ethical Approval Code No. LCWU: 2025:254).

Author Contributions:

Z. A., D. K., and H. A. Conceptualization, and Methodology. B. S. A., N. H., Data Curation, Formal Analysis and Investigation Z. S., R. H., and M. H. M., Writing – Original Draft Preparation. Z. S., and B. S. A., Project Administration, Validation, Writing, and Revision.

Declaration of Competing Interest:

The authors declare that they have no competing financial interests.

Funding

None

REFERENCES

Abalaka, S. E., Oyelowo, F. O., Akande, M. G., Tenuche, O. Z., Sani, N. A., Adeyemo, B. T., & Ejeh, S. A. (2021). Effects of *Moringa oleifera* leaves extract, vitamin C, and taurine co-exposures on calcium and metallothionein levels, oxidative stress, and gill histopathological changes in *Clarias gariepinus* exposed to sub-lethal cadmium. *Environmental Science and Pollution Research*, 28(37), 52258-52271. DOI: [10.1007/s11356-021-14426-z](https://doi.org/10.1007/s11356-021-14426-z)

Abdelwahab, A. M., El-Bahr, S. M., & Al-Khamees, S. (2020). Influence of dietary garlic (*Allium sativum*) and/or ascorbic acid on performance, feed utilization, body composition and hemato-biochemical parameters of juvenile Asian Sea Bass (*Lates calcarifer*). *Animals*, 10(12), 2396. DOI: [10.3390/ani10122396](https://doi.org/10.3390/ani10122396)

Abdul-Qadir, A. M., Aliyu-Paiko, M., Mohammed, A. K., & Ndejiko, M. J. (2024). Biofloc technology as a sustainable alternative for managing aquaculture wastewater. *Journal of Biochemistry, Microbiology and Biotechnology*, 12(1), 44-53. DOI: 10.1111/ire.12449

Alberts, A., Moldoveanu, E. T., Niculescu, A. G., & Grumezescu, A. M. (2025). Vitamin C: a comprehensive review of its role in health, disease prevention, and therapeutic potential. *Molecules*, 30(3), 748. DOI: 10.3390/molecules30030748

Awuchi, C. G., Igwe, V. S., Amagwula, I. O., & Echeta, C. K. (2020). Health benefits of micronutrients (vitamins and minerals) and their associated deficiency diseases: A systematic review. *International Journal of Food Sciences*, 3(1), 1-32. DOI: 10.47604/ijf.1024

Biswas, P., Dellanoce, C., Vezzoli, A., Mrakic-Sposta, S., Malnati, M., Beretta, A., & Accianni, R. (2020). Antioxidant activity with increased endogenous levels of vitamin C, E and A following dietary supplementation with a combination of glutathione and resveratrol precursors. *Nutrients*, 12(11), 3224. DOI: 10.3390/nu12113224

Daniel, N., Muralidhar, A. P., Srivastava, P. P., Jain, K. K., Prasad, K. P., & Ranjan, A. (2021). Effect of vitamin C on immune and stress responses in striped catfish, *Pangasianodon hypophthalmus* juveniles under pre-and post-challenge with *Aeromonas hydrophila*. *Aquaculture Research*, 52(12), 6444-6452. DOI: 10.1111/are.15509

El-Gendy, K. S., Aly, N. M., Mahmoud, F. H., Kenawy, A., & El-Sebae, A. K. H. (2010). The role of vitamin C as antioxidant in protection of oxidative stress induced by imidacloprid. *Food and chemical Toxicology*, 48(1), 215-221. DOI: 10.1016/j.fct.2009.10.003

Fasil, D. M., Hamdi, H., Al-Barty, A., Zaid, A. A., Parashar, S. K. S., & Das, B. (2021). Selenium and zinc oxide multinutrient supplementation enhanced growth performance in zebra fish by modulating oxidative stress and growth-related gene expression. *Frontiers in Bioengineering and Biotechnology*, 9, 721717. DOI: 10.3389/fbioe.2021.721717

Fazal, R. M., Fatima, A., Al Sulivany, B. S., Hussain, R., Hassan, A., Shahid, A., & Owais, M. (2025). Enhancing Growth Performance, Antioxidant Defense, And Immune Response in Striped Catfish (*Pangasius hypophthalmus*) Through Dietary Supplementation with Aloe Vera: A Sustainable Aquaculture Approach. *Science Journal of University of Zakho*, 13(4), 510-518. DOI: 10.25271/sjuz.2025.13.4.1691

Firouzi, S., Pahlavani, N., Navashenaq, J. G., Clayton, Z. S., Beigmohammadi, M. T., & Malekhamadi, M. (2022). The effect of Vitamin C and Zn supplementation on the immune system and clinical outcomes in COVID-19 patients. *Clinical nutrition open science*, 44, 144-154. DOI: 10.1016/j.nutos.2022.06.006

Gao, X., Wei, M., Ma, D., Yang, X., Zhang, Y., Zhou, X., & Yang, W. (2021). Engineering of a hollow-structured Cu2-XS nano-homojunction platform for near infrared-triggered infected wound healing and cancer therapy. *Advanced Functional Materials*, 31(52), 2106700. DOI: 10.1002/adfm.202106700

Gupta, A., Gupta, S. K., Priyam, M., Siddik, M. A., Kumar, N., Mishra, P. K., & Pattanayak, A. (2021). Immunomodulation by dietary supplements: A preventive health strategy for sustainable aquaculture of tropical freshwater fish, *Labeo rohita* (Hamilton, 1822). *Reviews in Aquaculture*, 13(4), 2364-2394. DOI: 10.1111/raq.12581

Hossain, M. A., Hossain, M. A., Haque, M. A., Mondol, M. M. R., Harun-Ur-Rashid, M., & Das, S. K. (2022). Determination of suitable stocking density for good aquaculture practice-based carp fattening in ponds under drought-prone areas of Bangladesh. *Aquaculture*, 547, 737485. DOI: 10.1016/j.aquaculture.2021.737485

Ibraheem, M., Mohamed, A. M. & Albassam. N. (2017). The Bacteria contamination of red local meat and white imported meat in Samaraa and Al-Dour districts. *Annals of Agricultural Science*, Moshtohor, 55(4), 831-836. DOI: 10.21608/assjm.2017.57017

Ibrahim, R. E., Ahmed, S. A., Amer, S. A., Al-Gabri, N. A., Ahmed, A. I., Abdel-Warith, A. W. A., & Metwally, A. E. (2020). Influence of vitamin C feed supplementation on the growth, antioxidant activity, immune status, tissue histomorphology, and disease resistance in Nile tilapia, *Oreochromis niloticus*. *Aquaculture Reports*, 18, 100545. DOI: 10.1016/j.aqrep.2020.100545

Ibrahim, R. E., Amer, S. A., Farroh, K. Y., Al-Gabri, N. A., Ahmed, A. I., El-Araby, D. A., & Ahmed, S. A. (2021). The effects of chitosan-vitamin C nanocomposite supplementation on the growth performance, antioxidant status, immune response, and disease resistance of Nile tilapia (*Oreochromis niloticus*) fingerlings. *Aquaculture*, 534, 736269. DOI: 10.1016/j.aquaculture.2020.736269

Ilhan, N., Kamanli, A., Ozmerdivenli, R., & Ilhan, N. (2004). Variable effects of exercise intensity on reduced glutathione, thiobarbituric acid reactive substance levels, and glucose concentration. *Archives of medical research*, 35(4), 294-300. DOI: 10.1016/j.arcmed.2004.03.006

Irshath, A. A., Rajan, A. P., Vimal, S., Prabhakaran, V. S., & Ganesan, R. (2023). Bacterial pathogenesis in various fish diseases: recent advances and specific challenges in vaccine development. *Vaccines*, 11(2), 470. DOI: 10.3390/vaccines11020470

Jiménez-Fernández, E., Ponce, M., Rodríguez-Rúa, A., Manchado, M., & Fernández-Díaz, C. (2018). Assessing the role of vitamin C and iron in early larvae stages of *Solea senegalensis* fed enriched Artemia. *Aquaculture*, 488, 145-154. DOI: 10.1016/j.aquaculture.2018.01.021

Katsoulos, P. D., Athanasiou, L. V., Karatzia, M. A., Giadinis, N., Karatzias, H., Boscos, C., & Polizopoulou, Z. S. (2017). Comparison of biuret and refractometry methods for the serum total proteins measurement in ruminants. *Veterinary Clinical Pathology*, 46(4), 620-624. DOI: 10.1111/vcp.12532

Liua, B., Wana, J., Gea, X., Xiea, J., Zhoub, Q., Miaob, L., & Panb, L. (2016). Effects of dietary Vitamin C on the physiological responses and disease resistance to pH stress and *Aeromonas hydrophila* infection of *Megalobrama amblycephala*. *Turkish Journal of Fisheries and Aquatic Sciences*, 16(2), 421-433. DOI: 10.4194/1303-2712-v16_2_22

Magouz, F. I., Dawood, M. A., Salem, M. F., El-Ghandour, M., Van Doan, H., & Mohamed, A. A. (2020). The role of a digestive enhancer in improving the growth performance, digestive enzymes activity, and health condition of Nile tilapia (*Oreochromis niloticus*) reared under suboptimal temperature. *Aquaculture*, 526, 735388. DOI:10.1016/j.aquaculture.2020.735388

Misra, C. K., Das, B. K., Mukherjee, S. C., & Pradhan, J. (2007). Effects of dietary vitamin C on immunity, growth and survival of Indian major carp *Labeo rohita*, fingerlings. *Aquaculture Nutrition*, 13(1), 35-44. DOI: 10.1111/j.1365-2095.2007.00451.x

Mohamad Hassan, S., Salah, M., Albassam, N. H., Mdloul, N. S., Muhaimeed, A. R. and Sulaiman, M. A. (2025). Using the Diverse Vegetables as a Filtration plants in Aquaculture Intensive System. *Tikrit Journal for Agricultural Sciences*, 25(1), 1-16. DOI: 10.25130/tjas.25.1.1

Moore, J. C., DeVries, J. W., Lipp, M., Griffiths, J. C., & Abernethy, D. R. (2010). Total protein methods and their potential utility to reduce the risk of food protein adulteration. *Comprehensive reviews in food science and food safety*, 9(4), 330-357. DOI: 10.1111/j.1541-4337.2010.00114.x

Namiq, K., Albassam, N. H., & Wahab, N. K. (2025). Assessment of Pesticide Residues in Fish Tissues and Water at Various Time of the Year. *Diyala Agricultural Sciences Journal*, 17(1), 178–193. DOI: 10.52951/dasj.25170114

Narra, M. R., Rajender, K., Reddy, R. R., Rao, J. V., & Begum, G. (2015). The role of vitamin C as antioxidant in protection of biochemical and haematological stress induced by chlorpyrifos in freshwater fish *Clarias batrachus*. *Chemosphere*, 132, 172-178. DOI: 10.1016/j.chemosphere.2015.03.006

Pavlou, S., Wang, L., Xu, H., & Chen, M. (2017). Higher phagocytic activity of thioglycollate-elicited peritoneal macrophages is related to metabolic status of the cells. *Journal of inflammation*, 14(1), 4. DOI: 10.1186/s12950-017-0151-x

Privat, C., Lantoine, F., Bedioui, F., Van Brussel, E. M., Devynck, J., & Devynck, M. A. (1997). Nitric oxide production by endothelial cells: comparison of three methods of quantification. *Life Sciences*, 61(12), 1193-1202. DOI: 10.1016/S0024-3205(97)00661-9

Rothe, G., Emmendorffer, A., Oser, A., Roesler, J., & Valet, G. (1991). Flow cytometric measurement of the respiratory burst activity of phagocytes using dihydrorhodamine 123. *J Immunol Methods*, 138(1), 133-5. DOI: 10.1016/0022-1759(91)90074-p

Sherif, A. H., Khalil, R. H., Talaat, T. S., Baromh, M. Z., & Elnagar, M. A. (2024). Dietary nanocomposite of vitamin C and vitamin E enhanced the performance of Nile tilapia. *Scientific Reports*, 14(1), 15648. DOI: 10.1038/s41598-024-65507-1

Singh, P., Kesharwani, R. K., & Keservani, R. K. (2017). Antioxidants and vitamins: Roles in cellular function and metabolism. In *Sustained energy for enhanced human functions and activity* 385-407. Academic Press.

Soldado, D., Bessa, R. J., & Jerónimo, E. (2021). Condensed tannins as antioxidants in ruminants Effectiveness and action mechanisms to improve animal antioxidant status and oxidative stability of products. *Animals*, 11(11), 3243. DOI: 10.3390/ani11113243

Surai, P. F., Kochish, I. I., Fisinin, V. I., & Kidd, M. T. (2019). Antioxidant defence systems and oxidative stress in poultry biology: An update. *Antioxidants*, 8(7), 235. DOI: 10.3390/antiox8070235

Taalab, H. A., Mohammady, E. Y., Hassan, T. M., Abdella, M. M., & Hassaan, M. S. (2022). β -Carotene of *Arthospira platensis* versus vitamin C and vitamin E as a feed supplement: Effects on growth, haemato-biochemical, immune-oxidative stress and related gene expression of Nile tilapia fingerlings. *Aquaculture Research*, 53(13), 4832-4846. DOI: 10.1111/are.15977

Tang, Z., Xie, S., Cui, Y., Zhan, W., Deng, Y., Peng, H., ... & Zhou, Q. (2024). Vitamin C as a functional enhancer in the non-specific immune defense, antioxidant capacity and resistance to low-temperature stress of juvenile mud crab, *Scylla paramamosain*. *Fish & Shellfish Immunology*, 153, 109834. DOI: 10.1016/j.fsi.2024.109834

Xu, C. M., Yu, H. R., Li, L. Y., Li, M., Qiu, X. Y., Fan, X. Q., ... & Shan, L. L. (2022). Effects of dietary vitamin C on the growth performance, biochemical parameters, and antioxidant activity of coho salmon *Oncorhynchus kisutch* (Walbaum, 1792) Postsmolts. *Aquaculture nutrition*, 2022(1), 6866578. DOI: 10.1155/2022/6866578

Xue, M., Wu, Y., Hong, Y., Meng, Y., Xu, C., Jiang, N., ... & Zhou, Y. (2022). Effects of dietary *Bacillus amyloliquefaciens* on the growth, immune responses, intestinal microbiota composition and disease resistance of yellow catfish, *Pelteobagrus fulvidraco*. *Frontiers in Cellular and Infection Microbiology*, 12, 1047351. DOI:10.3389/fcimb.2022.1047351

Zhang, A., Stacey, H. D., D'Agostino, M. R., Tugg, Y., Marzok, A., & Miller, M. S. (2023). Beyond neutralization: Fc-dependent antibody effector functions in SARS-CoV-2 infection. *Nature Reviews Immunology*, 23(6), 381-396. DOI: 10.1038/s41577-022-00813-1