

EVALUATION OF OXIDATION – ANTIOXIDATION BALANCE IN SERA OF PATIENTS WITH ACUTE HEPATITIS VIRUS TYPE C.

Rasha Hasan Jasim

Chemistry Department-College of Education for Girls-University of Kufa-Iraq

(Accepted for publication: June 9, 2013)

Abstract

Aim of the present study has to focused on the possible relationship between serum malondialdehyde level, an index of lipid peroxidation, and ceruloplasmin levels, as protective agent against lipid peroxidation, in hepatitis C virus. A group of 31 hepatitis virus type C patients were enrolled in the study, while; control group consisted of 25 healthy subjects. In the present study, total proteins (g/L), malondialdehyde (μM), ceruloplasmin oxidase activity (U/L) and ceruloplasmin concentration (g/L) were measured in sera samples of patients with hepatitis C virus as well as in the healthy controls.

Key words: lipid peroxidation, malondialdehyde, ceruloplasmin

Introduction

Viral hepatitis is the major health problem in Iraq, the most important causes of spreading the disease are low levels of socioeconomical status and poor hygiene conditions, particularly in countryside [Saravanamuttu, et al. 2007]. Acute viral hepatitis is one of the most common infectious diseases [Mustafa, et al. 2006], while hepatitis C virus (HCV) is the most baleful form of acute viral hepatitis in many parts of the world [Fujinaga, et al. 2011], Hepatitis C mostly occurs in the context of community-wide epidemics during which infection is transmitted from person to person by the fecal-oral route [Albert, et al. 2012].

Epidemically of HCV may habit to hypothesis that HCV is one of the main causative agents of chronic viral hepatitis. Chronic hepatitis C can progress to cirrhosis and eventually to hepatocellular carcinoma over a period of 20 to 30 years [Ulrich, et al. 2012]. The mechanisms which explain how HCV causes cell damage are not well understood, but symphoniously with the fact that liver is the central organ in the metabolism processes [Ghorbani, et al. 2008], one of the mechanisms including immunological liver damage, direct cytotoxicity mediated by different viral product and inductions of oxidative stress have been suggested as playing a pathogenic role in this infection, It has been suggested that HCV may cause oxidative stress in infected cell [Haimin, et al. 2006 and Görennek, et al. 2006].

Oxidative stress is defined as an imbalance between pro-oxidant and antioxidant mechanisms in our body, the balance being tilted in favor of the former, It has been unequivocally

implicated in the pathogenesis and pathophysiology of many diseases [Sumit, et al. 2008]. Oxidative stress is associated with the disturbance of hepatocyte biochemistry and generation of ROS (reactive oxygen species), when the antioxidant defenses are critically lowered [Mehmet, et al. 2004].

These ROS can damage cells by causing lipid peroxidation, and oxidative damage of DNA and proteins, and by depleting ATP stores. Prime targets of peroxidation by ROS are polyunsaturated fatty acids (PUFA) in membrane lipids. In the presence of metals (such as Fe^{3+} , Cu^{2+}), O^{2-} can react with H_2O_2 to generate a hydroxyl radical than become even more reactive and cytotoxic than O^{2-} or H_2O_2 [Görennek, et al. 2006]. PUFA is degraded by free radicals to form malondialdehyde (MDA). The level of MDA in serum serves as a marker of cellular damage due to free radicals. Cells have multiple mechanisms to remove free radicals and thereby minimize tissue injury. Antioxidants such as enzymes, like: superoxide dismutase (SOD) and Ceruloplasmin (Cp), catalase (like glutathione "GSH"), and nutritional antioxidants (vitamins E and C) trap free radicals and act as free radical scavenging systems [Nikam, et al. 2009, Sahu et al. 2009 and Moreno-Otero, et al. 2010].

Aim of the present study work to investigate the possible relationship between serum malondialdehyde level, an index of lipid peroxidation, and ceruloplasmin levels, as protective agents against lipid peroxidation, in acute hepatitis C virus cases.

Materials and Methods

Patients

The study group comprised 31 patients with newly diagnosed hepatitis C virus, with age range 23-67 years. They were admitted consecutively to the Liver and Digestive Tract Center of Al-Sader Medical City in Najaf, Iraq, between November 2010 and June 2011. All

patients were enrolled in the study **before receiving the course of drugs**. The patients group consisted of 17 females and 14 males and their average age was 44 years. The control group comprised 25 healthy individuals that included 12 females and 13 males aged between 19-62 (average of 43), the ratio of male to female was shown in **Figure 1**.




Figure 1: Gender Distribution of The Study Groups' Individuals

Blood Samples

All sera were collected in the morning after fasting 10 hour feature of the subjects in the present study are shown in the table 1. The healthy volunteers were selected on the basis of no alcoholic and smoking habits, without history of viral hepatitis, routing clinical check up during the entire period of research. Patients and control subjects residing in the same

geographical area, and they were in the same socioeconomic status and similar diet habits. Patients with chronic hepatitis C diagnosed based on clinical, biochemical, histological and virological evidence in the same medical city. Blood samples were taken from subjects in accordance with standard procedure, 5 ml of blood was collected from vein and protected in evacuated tubes without anticoagulation agents.

Table 1: Levels (g/L) of TSP, (μM) of MAD, (U/L) of Ceruloplasmin Oxidase Activity, and (g/L) of Ceruloplasmin Concentration in Patient of Hepatitis C Virus and control subjects (Mean \pm S.D.)

	Age	TSP Level	MAD	Cp. Activity	Cp. Level	Cu Level	Fe Level
	(Years)	(g/L)	(μM)	(U/L)	(g/L)	($\mu\text{g/ml}$)	($\mu\text{g/ml}$)
Individuals							
	Mean \pm S.D.	Mean \pm S.D.	Mean \pm S.D.	Mean \pm S.D.	Mean \pm S.D.	Mean \pm S.D.	Mean \pm S.D.
	Min-Max	Min-Max	Min-Max	Min-Max	Min-Max	Min-Max	Min-Max
Patients	42.521	66.344 \pm 8.710	20.534 \pm 7.241	15.173 \pm 7.882	4.233 \pm 2.521	0.422 \pm 0.114	1.937 \pm 0.891
(n=31)	\pm 15.701 23-67	53-81	9.375-28.513	8.263-23.991	3.535-9.530	0.133-0.753	1.071-3.181
Controls	41.922 \pm 15.213	74.122 \pm 9.095	9.581 \pm 2.632	43.024 \pm 10.338	8. 942 \pm 4.852	0.692 \pm 0.291	1.176 \pm 0.391
(n=25)	19-62	59-83	3.325-14.381	15.705-63.169	6.723- 15.089	0.147 - 1.112	0.900-1.483
p-value	0.583	0.064	0.001	0.000	0.001	0.001	0.000

Determination of Total Serum Proteins Levels

A total serum protein was estimated using Biuret method [Plummer 1971]. Biuret reagent supplied by the Manufacturing Company in a container contains 100 ml, that consist of sodium hydroxide (100 mM), sodium-potassium tartrate (16 mM), potassium iodide (15 mM), and cupric sulfate (6 mM). The bovine serum albumin was used as a standard protein. The procedure included mixing of 50 μ l serum or standard with 2.5 ml Biuret reagent, then the mixture was incubated at room temperature for 30 minutes, and the absorbance was measured at 540 nm.

Measurement of Serum Malondialdehyde Level

Malondialdehyde level is measured by the thiobarbituric acid-reacting substances (TBARS) assay [Yagi 1982]. Abridgment, 150 μ l of the serum sample was mixed with 1 ml of trichloro acetic acid (TCA) (17.5 %) and 1ml of thiobarbituric acid (0.6 %). The final mixture was mix using the vortex, then heated at 100°C for 15 minutes in the water bath. After the mixture was cooled with tap water, it was extracted with 1 ml TCA (70 %), the mixture was stand for 20 minutes at 25°C, and centrifuged at 3000 xg for 15 minutes. The organic phase was measured by use of a spectrophotometer with a wavelength of 534 nm.

Determination of Ceruloplasmin Oxidase activity

The activity of ceruloplasmin oxidase was determined in serum using the modified Rice method [Rice 1962]. The procedure included two glass tubes, test (A) and blank (B), 1ml of substrate buffer was added to each tube, then incubated at 37°C for 5 min. A 100 μ l of serum sample was added to tube A then incubated at 37°C for 15 min. A volume of 3 ml of cold working inhibition solution was added to all of A and B tubes; at last 100 μ l of deionized water was added to tube B. The absorbance was measured at $\lambda=540$ nm.

Determination of Serum Copper and Iron Levels

The levels of serum copper and iron were determined by flame atomic absorption spectrophotometry (GBC-933plus).

Statistical Analysis

The results were expressed as the mean \pm standard deviation (S.D.). The data were analyzed with **Student's independent t test**. All statistical analyses were performed with the Statistical Package for the Social Science (SPSS for Windows, Version 19.0). Pearson's correlation was applied to determined the relations among the laboratory parameters of the present study, significance was determined regression. A *p*-value of <0.01 was accepted as statistically significant.

Results and Discussion

Results of the current study, total proteins, malondialdehyde (μ M), ceruloplasmin oxidase activity (U/L) and ceruloplasmin concentration (g/L) are summarized in **Table 1**. The results shown that no significant variation (*p* =0.064) total serum proteins' levels in patients with hepatitis C virus when compared with those of healthy individuals. On the other hand, the statistical evaluation failed to exhibit significant variation (*p*<0.001) for serum malondialdehyde when patients of hepatitis C virus were compared with those of healthy controls. With same way, when the comparison was made for hepatitis C virus patients and control group, highly significant variations were found for ceruloplasmin oxidase activity and ceruloplasmin concentration. Statistical analysis demonstrated a significant (*p*<0.001) variation in copper levels in hepatitis C virus patient group when compared with those in controls. On the other hand, elevation in iron levels in hepatitis C virus patients group comparison to controls group was recorded, the lowest copper levels and highest iron levels were observed in oldest patients.

In order to find the possible relation between the oxidation and antioxidation processes through the infection with C virus of hepatitis, the correlation between malondialdehyde as a profile to the peroxidation process, and ceruloplasmin enzyme (ceruloplasmin oxidase activity) as an agent of antioxidation was studied.

Correlation coefficient showed the percentage of negative correlation (*r* = - 0.813 at *p*<0.000) between malodialehyde levels and celoplasmin oxidase activity in sera of pateints (**Figure 2 A**), while no such correlation was observed in healthy control group (**Figure 2 B**).




Figure 2: Correlation Coefficient Between Serum Malondialdehyde and Ceruloplasmin Oxidase Activity in A: Patients of Hepatitis C Virus, and B: Healthy Subjects

In the same way, a significant negative correlation ($r = 0.731$ at $p < 0.001$) was observed between levels of serum malonaldehyde and ceruloplasmin concentration in patients group (**Figure 3 A**). On the other hand, the healthy individuals group failed to illustrate same correlation between serum malonaldehyde levels and ceruloplasmin concentration was carried out (**Figure 3 B**).




Figure 3: Correlation Of Serum Malondialdehyde To The Ceruloplasmin Concentration In A: Patients of Hepatitis C Virus, and B: Healthy Individuals

Results of hepatitis C infection parameters which evaluated in the current study, largely depends upon the patient's age at infection and immune status as well as the level of hepatitis C replication. According to the previous studies, normally; total serum proteins level decrease

with the age progression; as well as this alteration (increase or decrease) in the protein levels was recorded in numerous diseases [Jasim 2009, Al-Faham 2002, Porter, et al. 2003, Raymond 2007, and Jasim 2011]. The result of current study agreed with these findings, where

the decreases in the total serum proteins' levels were proportion to the individuals' age (patients and controls), in addition to that the lowest proteins levels were recorded in sera of elderly individuals.

Oxidative damage by oxygen free radicals such as superoxide anions are known to be one of the factors involved in the mechanisms of diseases [Harliansyah, et al. 2005]. During immune activation by viral entities, neutrophils and others cells produce ROS as a mechanism of signal amplification for protection [Vera, et al. 2012]. ROS have also been implicated in a number of hepatic pathologies in exacerbating liver diseases [Jasim 2011], however, the oxidant production associated with immune reactions against viral hepatitis may lead to the formation of hepatocellular carcinoma, but unfortunately, approximately 10-40% of the patients with chronic HCV treat with the modern therapeutic models maintain the rate of sustained virologic response, therefore; detection of HCV infection and the effective advancement in the antiviral treatments against chronic HCV is necessary [Ko, et al. 2005].

In previous study (personality work in our laboratory from 2010-2012) [Jasim 2011, Harliansyah, et al. 2005, Vera, et al. 2012, Vera, et al. 2012, and Jasim 2011], it showed a progressive rise of malondialdehyde level in serum samples for individuals (patients and healthy) with age. This relationship was highly significant. In the present work, there was a significant negative correlation between serum malondialdehyde level and the age progression (data not shown); this result agreed with the previous findings. On the other hand; malonaldehyde level in sera of patients was elevated (100% of cases) than those in sera of healthy individuals; according to this observation the lipid peroxidation status occurred in acute and chronic of infection with HCV, when compared this results with the previous study that deal with sera of patients suffered infection with hepatitis virus type B, this study showed lipid peroxidation status didn't occurred in the primary stage of infection with hepatitis virus type B. **This comparison illustrates important finding, we can exploit oxidative stress parameter (MDA) in differentiation between hepatitis virus B and C types, to suggest the suitable therapy.** This result is agreed with several studies that measured malonaldehyde level as a reflex for lipid peroxidation in numerous diseases [27, 28],

while it disagreed with other studies, especially those with liver diseases [McGrath, et al. 2001, Aksoy, et al. 2003, Ko, et al. 2005, Janaia, et al. 2010, and Jasim 2011].

Ceruloplasmin is a blue alpha-2 glycoprotein with a molecular weight of 132,000u, It binds 90-95% of blood plasma copper, has 6-7 Cu ions per molecule, and exhibits ferroxidase activity (an iron oxidase), amine oxidase activity, superoxidase activity, as well as it is involved in Cu transport and homeostasis [Viorica, et al. 2009].

The symptoms of many diseases that is characterized by low levels of ceruloplasmin with subsequent copper deposition in various tissues [Merle, et al. 2010]. Results of present study agreed with these studies. In addition, ceruloplasmin is an acute phase reactant, whose concentration increases in inflammation, infection, and trauma, for these properties it is known as an antioxidant. This fact boost the observations of the current work which explained the decreases in Cp activity and concentration in HCV serum to the liver cell damage which causes countermarch in liver functions, then drop in synthesis of many proteins, ceruloplasmin may one of them.

Concurrent study designed in order to investigate the roles and alterations of ceruloplasmin in acute and chronic HCV and compare its outcome with previous study that carried out on the chronic HBV samples.

For the propose of arrive to integral understanding about the ceruloplasmin in HCV case, because copper and iron are linked to the ceruloplasmin [Patel, et al. 2002 and Brewer, et al. 2010], their levels were determined. Results of the current study illustrated that decrease of copper level in serum of patients with hepatitis virus type C comparison to healthy controls, adverse result was found at the levels of iron determined. Highlight on the results of copper and iron levels in the patients' sera illustrate that the decrease of copper concentration can be explained as a reflex to the decrease in the ceruloplasmin oxide activity and ceruloplasmin concentration as antioxidant agent in HCV, while, the literatures recorded that raises of the iron levels can be refer to the damage of liver cells [Aigner 2008, Bush 2010], these information support the hypothesis of the present work "liver cells may damage during different HCV infection stages".

In order to prove the hypothesis of liver cells damage during the prim and advanced infection

periods with hepatitis virus C type, the relationship of the malondialdehyde levels to ceruloplasmin oxidase activity, and ceruloplasmin concentration was carried out. The study outcome showed there are significant relation between these parameters; for that, we can conclude that a decrease in the ceruloplasmin oxide activity and ceruloplasmin concentration are reflex to arise in the lipid peroxidant products and an ambulance in the oxidation – antioxidation status.

References

- Aigner E and Datez C. [2008]: **Iron Perturbations in Human Non-Alcoholic Fatty Liver Disease (NAFLD): Clinical Relevance and Molecular Mechanisms.** *Hepatitis Monthly.* 8(3): 213-220.
- Aksoy H, Koruk M, and Akçay F. [2003]: **The Relationship Between Serum Malondialdehyde and Ceruloplasmin in Chronic Liver Disease.** *Turkish Journal of Biochemistry.* 28 (2): 32-34.
- Albert J C and Deborah K F.[2012]: **Diagnosis And Treatment Of Autoimmune Hepatitis.** *Hematology.* 36(2):479-497.
- Al-Faham R. [2002]: **A Biochemical Study of Serum Sialic Acid and Lectins Activity as Tumor Markers for Gastrointestinal Tract Cancers.** University of Kufa, College of Education, Chemistry Department. Thesis.
- Brewer G J, Kanzer S H, Zimmerman E A, Celmins D F, Heckman S M.[2010]: **Copper and Ceruloplasmin Abnormalities in Alzheimer's Disease.** *American Journal of Alzheimer's Disease.* 25 (6): 490-497.
- Bush A I. [2010]: **Iron-Export Ferroxidase Activity of β -Amyloid Precursor Protein Is Inhibited by Zinc in Alzheimer's Disease.** *Cell Journal.* 142(6): 857-867.
- Fujinaga H, Tsutsumi T, Yotsuyanagi H, Moriya K, And Koike K. [2011]: **Hepatocarcinogenesis In Hepatitis C: HCV Shrewdly Exacerbates Oxidative Stress By Modulating Both Production And Scavenging Of Reactive Oxygen Species.** *Oncology.* 81(1):11-17.
- Ghorbani G A, Alavian S M, and Ghadimi H R. [2008]: **Long Term Effect of One or Two Doses of Hepatitis C Vaccine in Adult after Five Years.** *Pakistan Journal of Biological Sciences.* 11(4): 660-663.
- Görenek L, Acar A, Aydın A, Eyigun C P, Çetinkaya A, Eken A, and Sayal A. [2006]: **Oxidative Stress And Antioxidant Defense In Patients With Chronic Hepatitis C Patients Before And After Pegylated Interferon Alfa-2b Plus Ribavirin Therapy.** *Journal of Translational Medicine.* 4(25): 1-6.
- Haimin C, Xiaojun Y, Peng Z, and Jing L. [2006]: **Antioxidant Activity And Hepatoprotective Potential Of Agar-Oligosaccharides In Vitro And In Vivo.** *Nutrition Journal.* 5(3): 1-12.
- Harliansyah A H, Noor A M, Wan Z, and Yasmin A M. [2005]: **Effects of Zingiber officinale on Superoxide Dismutase, Glutathione Peroxidase, Catalase, Glutathione and Malondialdehyde Content in HepG2 Cell Line.** *Malaysian Journal of Biochemistry and Molecular Biology.* 11, 36-41.
- Janaia A V and Suprapaneni K M. [2010]: **Antioxidant Vitamins And Enzymes Status In Patients With Alcoholic Liver Disease.** *Journal of Clinical and Diagnostic Research.* 4(4): 2742-2747.
- Jasim R H. [2009]: **Isolation, Purification, and Partial Characterization of Human Lectins from Patients of Kidney Tumors.** Baghdad University, College of Science for Women, Chemistry Department. Treatise.
- Jasim R H. [2011]: **Serum Malondialdehyde And Lipid Profile Levels Of Young Patients "Haven't A Family History Of Hypertension": A New Study For Cases In The Civic Society.** *Pakistan Journal Of Chemistry.*1(3): 120-125.
- Jasim R H. [2011]: **Evaluation of Oxidative-Antioxidative Balance in Serum of Patients With Non Acute Hepatitis Virus Type B.** Under publishing (Participated in the 2011 International Conference on Chemistry and Chemical Process – ICCCP 2011)
- Ko W, Guo C, Yeh M, Lin L, Hsu G W, Chen P, Luo M, and Lin C. [2005]: **Blood Micronutrient, Oxidative Stress, and Viral Load in Patients with Chronic Hepatitis C.** *World Journal of Gastroenterology.* 11(30):4697-4702.
- McGrath L, Mc Gleenon B, Brennan S, McColl D, McIlroy S, and Passmore A. [2001]: **Increased Oxidative Stress in Alzheimer's Disease as Assessed with 4-**

- Hydroxynonenal but not Malondialdehyde.** Q J Med. 94: 485-490.
- Mehmet K, Seyithan T, Cemil S, Omer Y, Fatih A, and Metin K. [2004]: **Oxidative Stress and Enzymatic Antioxidant Status in Patients with Nonalcoholic Steatohepatitis.** Annals of Clinical and Laboratory Science. 34(1): 57-62.
- Merle U, Weiss K H, Eisenbach C, Tuma S, Ferenci P, and Stremmel W. [2010]: **Truncating Mutations in the Wilson Disease Gene ATP7B are Associated with Very Low Serum Ceruloplasmin Oxidase Activity and An Early Onset of Wilson Disease.** BMC Gastroenterology. 10(8): 2-6.
- Moreno-Otero R and Trapero-Marugán M. [2010]: **Hepatoprotective Effects of Antioxidants in Chronic Hepatitis C.** World Journal of Gastroenterol.16(15): 1937-1938 .
- Mustafa C, Semiha D, Fahri B, Hüseyin Ç, Fatma C, and Nihat M. [2006]: **Relationship Between Antioxidant Capacity And Oxidative Stress In Children With Acute Hepatitis A.** World Journal of Gastroenterology. 12(38): 6212-6215.
- Nikam S, Nikam P, Ahaley S K. [2009]: **Role of Free Radical and Antioxidant Imbalance in Pathogenesis of Parkinson's Disease.** Biomedical Research. 20 (1): 55-58.
- Patel B N, Dunn R J, Jeong S Y, Zhu Q, Julien J, and David S. [2002]: **Ceruloplasmin Regulates Iron Levels in the CNS and Prevents Free Radical Injury.** The Journal of Neuroscience. 22(15): 6578-6586.
- Plummer A. [1971]: **An Introduction to Practical Biochemistry.** 1st Edition. Mcgraw Hill. Book Company. London. U K.
- Porter G T, Pou A M, Quinn F B, & Ryan M W. [2003]: **Nutritional Considerations for Head and Neck Cancer Patients.** Grand Rounds Presentation, UTMB, Department of Otolaryngology.
- Raymond R W. [2007]: **Biochemistry of Cancer.** In: **Cancer Biology.** Section 1.p 108-120.
- Rice E W. [1962]: **Ceruloplasmin Assay in Serum: Standardization of Ceruloplasmin Activity in Terms of International Enzyme Unit "Standard Methods of Clinical Chemistry".** 4th Edition, Siligson D., New York, Academic press.
- Sahu S, Abraham R, Vedavallir R, and Daniel M. [2009]: **Study of Lipid Profile, Lipid Peroxidation, and Vitamin E in Pregnancy Induced Hypertension.** Indian Journal of Physiology Pharmacology. 53 (4): 365–369.
- Saravanamuttu G I, Samreen M J, Ramsay M, Jonathan G.[2007]: **International Public Health Repository for Hepatitis C.** Nucleic Acid Research. 35(2): 367-370.
- Sumit B, Mukul P A, Dwivedi S, and Banerjee BD. [2008]: **Monitoring Oxidative Stress Across Worsening Child Pugh Class Of Cirrhosis.** Indian Journal of Medical Science. 62(11): 444-451.
- Ulrich S. [2012]: **Diagnosis, Prognosis and Therapy of Hepatocellular Carcinoma.in heamatology.** Flying Publisher.UK.
- Vera G V, Goran G, Senijo P, Antonio S, Ana-Marija D, Dubravka F, Maja P. [2012]: **Assessment of cytogenetic damage and oxidative stress in personnel occupationally exposed to the pulsed microwave radiation of marine radar equipment.** International Journal of Hygiene and Environmental Health. 38(2): 1438-1449.
- Viorica L A, William H R, and Kirk L. [2009]: **Ceruloplasmin levels in human sera from various diseases and their correlation with patient's age and gender.** Health. 1(2),104-110.
- Yagi K. [1982]: **Assay for Lipid Peroxide Level and Its Clinical Significance.** In: **Yagi K Editor Lipid Peroxides in Biology and Medicine,** New York. p233–242.

الخلاصة

تهدف الدراسة الحالية الى امكانية ايجاد علاقة بين مستوى المألونالديهيد كدالة لحدوث حالة مافوق الاكسدة للدهون ومستوى السيرلوبيلازمين كعامل مضاد لحالة ما فوق الاكسدة للدهون في امصال مرضى التهاب الكبد الفيروسي من النوع سي. ضمت الدراسة 31 مصاباً بالتهاب الكبد الفيروسي من النوع سي كعينة للدراسة بينما ضمت مجموعة السيطرة 25 شخصاً سليماً. في الدراسة الحالية تم تقييم مستويات كل من البروتينات الكلية (غم/ليتر) والمألونالديهيد (مايكرمولا ري) وفعالية السيرلوبيلازمين المؤكسد (وحدة انزيمية/ليتر) و تركيز السيرلوبيلازمين (غم/ليتر) في العينات المصلية لكل من مرضى التهاب الكبد الفيروسي من النوع سي و افراد مجموعة السيطرة.