

EXISTENCE, UNIQUENESS AND STABILITY OF PERIODIC SOLUTION FOR NONLINEAR SYSTEM OF INTEGRO-DIFFERENTIAL EQUATIONS

Raad Nori. Butris^{a,*}, Ava Shafeeq Rafeeq^b, Hewa Selman Faris^c^{a, c} Department of Mathematics, College of Basic Education, University of Duhok, Kurdistan Region, Iraq - (raad.butris@uod.ac, hewa.faris@uod.ac)^b Department of Mathematics, Faculty of Science, University of Zakho, Kurdistan Region, Iraq - ava.rafeeq@uoz.edu.krd**Received: Aug. 2016 / Accepted: Feb. 2017 / Published: Mar. 2017**<https://doi.org/10.25271/2017.5.1.312>**ABSTRACT:**

In this paper, we investigate the existence, uniqueness and stability of the periodic solution for the system of nonlinear integro-differential equations by using the numerical-analytic methods for investigate the solutions and the periodic solutions of ordinary differential equations, which are given by A. Samoilenko.

KEYWORDS: Existence, Uniqueness & Stability solution; Periodic Solution; Nonlinear System of Integro-differential Equations, Numerical-Analytic Methods.

1. INTRODUCTION

Analysis of various processes in mechanics, physics, biology and celestial mechanics, and other branches of science and engineering requires investigation into the existence and uniqueness solutions of differential equations of various types and systems (Apostol, 1973; Burrill, 1969). In particular, there developed many problems of the existence of periodic solutions and construction business. Among them, one should mention especially numerical-analytical method of successive approximations (Samoilenko, 1965; Tidke & More, 2015).

The conditions of existence solutions that are important in the theories of analysis and results are often obtained by using fixed-point theorems (Banach, Schauder) and successive approximation (Hu & Li, 2005; Rama, 1981). The study theories about the existence, uniqueness and stability of differential equations (David & Williams, 2015; Korol, 2010). There are many details to the investigation of different aspects of this theory (see, for example, (Butris et al., 2015; Byszewski et al., 1997; Korol, 2009). Previously, Samoilenko in many original works referred to as the numerical-analytic method based upon successive approximations (Samoilenko, 1965; Samoilenko, 1966). Tidke and More (Tidke & More, 2015), investigated the existence and uniqueness of solution of Volterra integro-differential equation with nonlocal condition in cone metric space. The result is obtained by using some extensions of Banach's contraction principle in complete cone metric space of the form.

$$x'(t) = A(t)x(t) + f(t, x(t)) + \int_0^t k(s, x(s))ds, \quad t \in J = [0, b] \text{ with } x(0) + g(x) = x_0 \quad \dots \quad (P1)$$

In (Aziz, 2006) Aziz, studied the periodic solutions for some systems of nonlinear ordinary differential equations which has the form:

$$\left. \begin{aligned} \frac{dx(t)}{dt} &= (A + B(t))x + f(t, x, y) \\ \frac{dy(t)}{dt} &= (C + D(t))y + g(t, x, y) \end{aligned} \right\} \quad (P2)$$

In this work, we study the existence, uniqueness and stability of solution of nonlinear system of integro-differential equations which has the form:-

$$\left. \begin{aligned} \frac{dx}{dt} &= Ax + f(t, \varepsilon, \int_{-\infty}^t R(t-\tau)(x(\tau) - y(\tau))d\tau) \\ \frac{dy}{dt} &= By + g(t, \mu, \int_{-\infty}^t G(t-\tau)(x(\tau) - y(\tau))d\tau) \end{aligned} \right\}, \quad (1.1)$$

and also investigate the existence and uniqueness of periodic solution of (1.1), where $\varepsilon = \int_a^b \lambda(x(\tau) - y(\tau))d\tau$, $\mu = \int_c^d \gamma(x(\tau) - y(\tau))d\tau$, λ and γ are constants, and $x \in D \subset \mathbb{R}^n$; $y \in D_1 \subset \mathbb{R}^n$, D and D_1 are closed and bounded domain. Assume that the vector functions $f(t, \varepsilon, u)$ and $g(t, \mu, v)$ where $u = \int_{-\infty}^t R(t-\tau)(x(\tau) - y(\tau))d\tau$ and $v = \int_{-\infty}^t G(t-\tau)(x(\tau) - y(\tau))d\tau$, are defined on the domain.

$$[t, x, y] \in \mathbb{R}^1 \times D \times D_1, \quad (1.2)$$

and continuous functions in t, x, y and satisfy the following inequalities:-

$$\left. \begin{aligned} \|f(t, \varepsilon, u)\| &\leq M \\ \|g(t, \mu, v)\| &\leq N \\ \|f(t, \varepsilon_1, u_1) - f(t, \varepsilon_2, u_2)\| &\leq j_1 \|\varepsilon_1 - \varepsilon_2\| + j_2 \|u_1 - u_2\| \\ \|g(t, \mu_1, v_1) - g(t, \mu_2, v_2)\| &\leq k_1 \|\mu_1 - \mu_2\| + k_2 \|v_1 - v_2\| \end{aligned} \right\} \quad (1.3)$$

for all $t \in \mathbb{R}^1$, $x \in D$, $y \in D_1$, where M, N, j_1, j_2, k_1 , and k_2 are positive constants. Suppose that the positive matrices A and B satisfy the condition:

$$\left. \begin{aligned} \|e^{A(t-\tau)}\| &\leq Q_1 \\ \|e^{B(t-\tau)}\| &\leq Q_2 \end{aligned} \right\} \quad (1.4)$$

* Corresponding author

where Q_1 and Q_2 are positive constant. Also, the functions $R(t - \tau)$ and $G(t - \tau)$ satisfying conditions

$$\left. \begin{aligned} \|R(t - \tau)\| &\leq \frac{he^{-\alpha(t-\tau)}}{(t-\tau)^{1-\alpha}} \\ \|G(t - s)\| &\leq \frac{\sigma e^{-\beta(t-\tau)}}{(t-\tau)^{1-\beta}} \end{aligned} \right\} \quad (1.5)$$

where $-\infty \leq 0 \leq \tau \leq t \leq T \leq \infty, 0 \leq \alpha \leq 1, 0 \leq \beta \leq 1, h$ and σ are positive constants, and the integrals of $\|R(t - \tau)\|$ and $\|G(t - \tau)\|$, and by using gamma function,

we can find that

$$\int_{-\infty}^t \|R(t - \tau)\| d\tau = p_\alpha \text{ and } \int_{-\infty}^t \|G(t - s)\| d\tau = p_\beta$$

where $p_\alpha = \alpha^{1-\alpha} h \Gamma(\alpha)$ and $p_\beta = \beta^{1-\beta} \sigma \Gamma(\beta)$.

The nonempty sets are as follow:-

$$\left. \begin{aligned} D_f &= D_0 - (I_1 + TQ_1M) \\ D_g &= D_1 - (I_2 + TQ_2N) \end{aligned} \right\} \quad (1.6)$$

where $I_1 = \|x_0\|(\|e^{At}\| + \|E\|)$ and $I_2 = \|y_0\|(\|e^{Bt}\| + \|E\|)$

Moreover, assume that the maximum eigen-value of the

matrix $\Lambda = \begin{pmatrix} TQ_1p_\alpha^\lambda & TQ_1p_\alpha^\lambda \\ TQ_2p_\beta^\gamma & TQ_2p_\beta^\gamma \end{pmatrix}$ is less than unity, i.e.

$$TQ_1p_\alpha^\lambda + TQ_2p_\beta^\gamma < 1. \quad (1.7)$$

2. EXISTENCE, UNIQUENESS AND STABILITY SOLUTION OF THE SYSTEM (1.1)

Theorem 2.1 (Existence theorem). Let the system (1.1) satisfy the inequalities (1.3)-(1.5) and the condition (1.6) and (1.7). Then the sequences of vectors functions $\{x_m(t)\}_{m=0}^\infty$ and $\{y_m(t)\}_{m=0}^\infty$ are defined by the following:

$$\begin{aligned} x_{m+1}(t, x_0, y_0) &= x_0 e^{At} \\ &+ \int_0^t e^{A(t-\tau)} f(t, \varepsilon_{m+1}, u_{m+1}) d\tau, x(0) \\ &= x_0, m = 0, 1, 2, \dots \end{aligned} \quad (2.1)$$

and,

$$\begin{aligned} y_{m+1}(t, x_0, y_0) &= y_0 e^{Bt} \\ &+ \int_0^t e^{B(t-\tau)} g(t, \mu_{m+1}, v_{m+1}) d\tau, y(0) \\ &= y_0, m = 0, 1, 2, \dots \end{aligned} \quad (2.2)$$

where $\varepsilon_{m+1} = \int_a^b \lambda(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0)) d\tau,$

$$\begin{aligned} \mu_{m+1} &= \int_c^d \gamma(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0)) d\tau, \\ u_{m+1} &= \int_{-\infty}^t R(t - \tau)(x_m(\tau, x_0, y_0) - \\ &y_m(\tau, x_0, y_0)) d\tau, v_{m+1} = \int_{-\infty}^t G(t - \tau)(x_m(\tau, x_0, y_0) - \\ &y_m(\tau, x_0, y_0)) d\tau, \end{aligned}$$

convergent uniformly when $m \rightarrow \infty$ in the domain (1.2) to the limit vectors functions $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ which are satisfying the following integral equations:-

$$\begin{aligned} x(t, x_0, y_0) &= x_0 e^{At} \\ &+ \int_0^t e^{A(t-\tau)} f(t, \varepsilon, u) d\tau, x(0) = x_0 \end{aligned} \quad (2.3)$$

$$\begin{aligned} y(t, x_0, y_0) &= y_0 e^{Bt} \\ &+ \int_0^t e^{B(t-\tau)} g(t, \mu, v) d\tau, y(0) = y_0 \end{aligned} \quad (2.4)$$

where $\varepsilon = \int_a^b \lambda(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau,$

$$\begin{aligned} \mu &= \int_c^d \gamma(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau, \\ u &= \int_{-\infty}^t R(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau, \\ v &= \int_{-\infty}^t G(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau, \end{aligned}$$

and it's a unique solution to the system (1.1), provided that

$$\left(\begin{aligned} \|x^0(t, x_0, y_0) - x_m(t, x_0, y_0)\| \\ \|y^0(t, x_0, y_0) - y_m(t, x_0, y_0)\| \end{aligned} \right) \leq \Lambda^m (E - \Lambda)^{-1} W_0 \quad (2.5)$$

where $W_0 = \begin{pmatrix} I_1 + TQ_1M \\ I_2 + TQ_2N \end{pmatrix},$

$\Lambda^m = \begin{pmatrix} TQ_1p_\alpha^\lambda & TQ_1p_\alpha^\lambda \\ TQ_2p_\beta^\gamma & TQ_2p_\beta^\gamma \end{pmatrix}^m,$ E is an identity matrix, for all $m \geq 1$ and $t \in R^1.$

Proof: From (2.1) and (2.2) when $m=0,$ we find that:

$$\begin{aligned} \|x_1(t, x_0, y_0) - x_0\| &\leq I_1 + TQ_1M, \text{ for all } t \in R^1, x_0 \in D_f. \\ \|y_1(t, x_0, y_0) - y_0\| &\leq I_2 + TQ_2N, \text{ for all } t \in R^1, y_0 \in D_g. \end{aligned}$$

By mathematical induction and from (2.1) and (2.2), we get:-

$$\|x_m(t, x_0, y_0) - x_0\| \leq I_1 + TQ_1M \quad (2.6)$$

$$\|y_m(t, x_0, y_0) - y_0\| \leq I_2 + TQ_2N \quad (2.7)$$

i. e. $x_m(t) \in D_0, y_m(t) \in D_1, x_0 \in D_f, y_0 \in D_g$ for all $t \in R^1.$

After that, we shall prove the sequences of vectors functions from (2.1) and (2.2) uniformly converges on domain(1.2), when $m > 1,$ we have

$$\begin{aligned} \|x_2(t, x_0, y_0) - x_1(t, x_0, y_0)\| \\ \leq TQ_1p_\alpha^\lambda (\|x_1(t, x_0, y_0) - x_0\| + \|y_1(t, x_0, y_0) - y_0\|) \end{aligned} \quad (2.8)$$

where $p_\alpha^\lambda = j_1 \lambda(b - a) + j_2 p_\alpha$

Similarly

$$\begin{aligned} \|y_2(t, x_0, y_0) - y_1(t, x_0, y_0)\| \\ \leq TQ_2p_\beta^\gamma (\|x_1(t, x_0, y_0) - x_0\| + \|y_1(t, x_0, y_0) - y_0\|) \end{aligned} \quad (2.9)$$

Where $p_\beta^\gamma = k_1 \gamma(d - c) + k_2 p_\beta$

By mathematical induction, we can get:-

$$\begin{aligned} \|x_{m+1}(t, x_0, y_0) - x_m(t, x_0, y_0)\| \\ \leq TQ_1p_\alpha^\lambda (\|x_m(t, x_0, y_0) - \\ x_{m-1}(t, x_0, y_0)\| \\ + \|y_m(t, x_0, y_0) - \\ y_{m-1}(t, x_0, y_0)\|) \end{aligned} \quad (2.10)$$

and

$$\begin{aligned} \|y_{m+1}(t, x_0, y_0) - y_m(t, x_0, y_0)\| \\ \leq TQ_2p_\beta^\gamma (\|x_m(t, x_0, y_0) - \\ x_{m-1}(t, x_0, y_0)\| \\ + \|y_m(t, x_0, y_0) - \\ y_{m-1}(t, x_0, y_0)\|) \end{aligned} \quad (2.11)$$

We rewrite the inequalities (2.10) and (2.11) in vector form, we find that

$$W_{m+1}(t) \leq \Lambda W_m(t) \quad (2.12)$$

where $W_{m+1}(t) = \begin{pmatrix} \|x_{m+1}(t, x_0, y_0) - x_m(t, x_0, y_0)\| \\ \|y_{m+1}(t, x_0, y_0) - y_m(t, x_0, y_0)\| \end{pmatrix},$

$W_m(t) = \begin{pmatrix} \|x_m(t, x_0, y_0) - x_{m-1}(t, x_0, y_0)\| \\ \|y_m(t, x_0, y_0) - y_{m-1}(t, x_0, y_0)\| \end{pmatrix},$ and

$$\Lambda = \begin{pmatrix} TQ_1 p_\alpha^\lambda & TQ_1 p_\alpha^\lambda \\ TQ_2 p_\beta^\gamma & TQ_2 p_\beta^\gamma \end{pmatrix}$$

Also from (2.12). We get

$$W_{m+1} \leq \Lambda^m W_0 \tag{2.13}$$

and hence

$$\sum_{j=1}^m W_j \leq \sum_{j=1}^m \Lambda^{j-1} W_0 \tag{2.14}$$

By condition (1.7), the inequality (2.14) is uniformly convergent when, $m \rightarrow \infty$, and

$$\lim_{m \rightarrow \infty} \sum_{j=1}^m \Lambda^{j-1} W_0 = \sum_{j=1}^{\infty} \Lambda^{j-1} W_0 = (E - \Lambda)^{-1} W_0 \tag{2.15}$$

Thus the sequence of vectors functions (2.1) and (2.1) are uniformly convergent. Suppose that

$$\left. \begin{aligned} \lim_{m \rightarrow \infty} x_m(t, x_0, y_0) &= x^0(t, x_0, y_0) \\ \lim_{m \rightarrow \infty} y_m(t, x_0, y_0) &= y^0(t, x_0, y_0) \end{aligned} \right\} \tag{2.16}$$

Since the sequences of function (2.1) and (2.2) are defined and continuous in the domain, $(t, x, y) \in [0, T] \times D \times D$, then the limiting functions $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ are also defined and continuous in the same domain.

Moreover, by (2.15) and proceeding (2.1) and (2.2) to the limit when $m \rightarrow \infty$ this show that the limiting functions $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ are the solutions of the integral equations (2.3) and (2.4).

Finally, we have to show that the solutions of system (1.1) are unique, suppose that $\tilde{x}(t, x_0, y_0)$ and $\tilde{y}(t, x_0, y_0)$ be another solution to the system (1.1), then

$$\tilde{x}(t, x_0, y_0) = x_0 e^{At} + \int_0^t e^{A(t-\tau)} f(t, \tilde{x}, \tilde{y}) d\tau, \tag{2.17}$$

$$\tilde{y}(t, x_0, y_0) = y_0 e^{Bt} + \int_0^t e^{B(t-\tau)} g(t, \tilde{x}, \tilde{y}) d\tau, \tag{2.18}$$

where

$$\begin{aligned} \tilde{\varepsilon} &= \int_a^b \lambda(\tilde{x}(\tau, x_0, y_0) - \tilde{y}(\tau, x_0, y_0)) d\tau; \tilde{u} \\ &= \int_{-\infty}^t R(t - \tau)(\tilde{x}(\tau, x_0, y_0) \\ &\quad - \tilde{y}(\tau, x_0, y_0)) d\tau, \end{aligned}$$

$$\begin{aligned} \tilde{\mu} &= \int_c^d \gamma(\tilde{x}(\tau, x_0, y_0) - \tilde{y}(\tau, x_0, y_0)) d\tau; \tilde{v} \\ &= \int_{-\infty}^t G(t - \tau)(\tilde{x}(\tau, x_0, y_0) \\ &\quad - \tilde{y}(\tau, x_0, y_0)) d\tau, \end{aligned}$$

Then,

$$\begin{aligned} \|x(t, x_0, y_0) - \tilde{x}(t, x_0, y_0)\| \\ \leq TQ_1 p_\alpha^\lambda (\|x(t, x_0, y_0) - \tilde{x}(t, x_0, y_0)\| \\ + \|y(t, x_0, y_0) - \tilde{y}(t, x_0, y_0)\|) \end{aligned} \tag{2.19}$$

$$\begin{aligned} \|y(t, x_0, y_0) - \tilde{y}(t, x_0, y_0)\| \\ \leq TQ_2 p_\beta^\gamma (\|x(t, x_0, y_0) - \tilde{x}(t, x_0, y_0)\| \\ + \|y(t, x_0, y_0) - \tilde{y}(t, x_0, y_0)\|) \end{aligned} \tag{2.20}$$

Rewrite the inequalities (2.19) and (2.20) in a vector form, we get

$$\begin{aligned} &\left(\|x(t, x_0, y_0) - \tilde{x}(t, x_0, y_0)\| \right. \\ &\quad \left. \|y(t, x_0, y_0) - \tilde{y}(t, x_0, y_0)\| \right) \\ &\leq \Lambda \left(\|x(t, x_0, y_0) - \tilde{x}(t, x_0, y_0)\| \right. \\ &\quad \left. \|y(t, x_0, y_0) - \tilde{y}(t, x_0, y_0)\| \right). \end{aligned} \tag{2.21}$$

where $\Lambda = \begin{pmatrix} TQ_1 p_\alpha^\lambda & TQ_1 p_\alpha^\lambda \\ TQ_2 p_\beta^\gamma & TQ_2 p_\beta^\gamma \end{pmatrix}$

Also from (2.21), we have:

$$\begin{aligned} &\left(\|x(t, x_0, y_0) - \tilde{x}(t, x_0, y_0)\| \right. \\ &\quad \left. \|y(t, x_0, y_0) - \tilde{y}(t, x_0, y_0)\| \right) \\ &\leq \Lambda^m \left(\|x(t, x_0, y_0) - \tilde{x}(t, x_0, y_0)\| \right. \\ &\quad \left. \|y(t, x_0, y_0) - \tilde{y}(t, x_0, y_0)\| \right). \end{aligned} \tag{2.22}$$

From the condition(1.7), we conclude that $x(t, x_0, y_0) = \tilde{x}(t, x_0, y_0)$ and $y(t, x_0, y_0) = \tilde{y}(t, x_0, y_0)$, then the solutions of (1.1) are unique. Thus the proof is complete.

Theorem 2.2. Let $x(t, x_0, y_0)$ and $y(t, x_0, y_0)$ are solutions of the system (1.1), and all condition of the theorem 2.1 are satisfied, then the following inequalities holds

$$\begin{aligned} \|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ \leq Q_1 F_1 F_3 \|x_0^1 - x_0^2\| \\ + TQ_1 Q_1 p_\alpha^\lambda F_1 F_2 F_3 \|y_0^1 - y_0^2\| \end{aligned} \tag{2.23}$$

$$\begin{aligned} \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\| \\ \leq TQ_1 Q_1 p_\beta^\gamma F_1 F_2 F_3 \|x_0^1 - x_0^2\| \\ + Q_2 F_2 F_3 \|y_0^1 - y_0^2\| \end{aligned} \tag{2.24}$$

for all $t \in R^1$, $x_0^1, x_0^2 \in D_f$ and $y_0^1, y_0^2 \in D_g$ where $F_1 = (1 - TQ_1 p_\alpha^\lambda)^{-1}$, $F_2 = (1 - TQ_2 p_\beta^\gamma)^{-1}$ and $F_3 = (1 - T^2 Q_1 Q_1 p_\alpha^\lambda p_\beta^\gamma F_1 F_2)^{-1}$.

Proof: Let $x(t, x_0^1, y_0^1)$, $y(t, x_0^1, y_0^1)$ and $x(t, x_0^2, y_0^2)$, $y(t, x_0^2, y_0^2)$ are two solutions of the system (1.1), then

$$\begin{aligned} \|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ \leq \|x_0^1 - x_0^2\| Q_1 \\ + TQ_1 p_\alpha^\lambda (\|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ + \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\|) \end{aligned} \tag{2.25}$$

Or

$$\begin{aligned} \|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ \leq \|x_0^1 - x_0^2\| Q_1 F_1 \\ + TQ_1 p_\alpha^\lambda F_1 \|y(t, x_0^1, y_0^1) \\ - y(t, x_0^2, y_0^2)\| \end{aligned} \tag{2.26}$$

and

$$\begin{aligned} \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\| \\ \leq \|y_0^1 - y_0^2\| Q_2 \\ + TQ_2 p_\beta^\gamma (\|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ + \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\|) \end{aligned} \tag{2.27}$$

or

$$\begin{aligned} \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\| \\ \leq \|y_0^1 - y_0^2\| Q_2 F_2 \\ + TQ_2 p_\beta^\gamma F_2 \|x(t, x_0^1, y_0^1) \\ - x(t, x_0^2, y_0^2)\| \end{aligned} \tag{2.28}$$

Substitute (2.28) in (2.26), we get (2.23) and substitute (2.26) in (2.28), we get (2.24). Hence, the proof is complete.

3. EXISTENCE, UNIQUENESS AND STABILITY OF PERIODIC SOLUTION OF THE SYSTEM (1.1)

In this section, we investigate the existence, uniqueness and stability of periodic solution of the problem of nonlinear system of differential equation (1.1).

Let the vector functions $f(t, \varepsilon, u)$ and $g(t, \mu, v)$ in (1.1) are defined on the domain

$$(t, x_0, y_0) \in [0, T] \times D_2 \times D_3, \tag{3.1}$$

where D_2 and D_3 are closed and bounded subsets of Euclidean spaces and $u = \int_{-\infty}^t R(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0))d\tau$, and $v = \int_{-\infty}^t G(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0))d\tau$, continuous functions in t, x, y and periodic in t of period T , i.e. $f(t, \varepsilon, u) = f(t + T, \varepsilon, u)$ and $g(t, \mu, v) = g(t + T, \mu, v)$. Also the vector functions $f(t, \varepsilon, u)$ and $g(t, \mu, v)$ satisfy (1.3) for all $t \in [0, T]$, and the matrices A and B satisfy the inequality (1.4), where $-\infty \leq 0 \leq \tau \leq t \leq T \leq \infty$.

We define the non-empty sets:

$$\left. \begin{aligned} D_{1f} &= D_2 - Q_1 T M \\ D_{1g} &= D_3 - Q_2 T N \end{aligned} \right\} \tag{3.2}$$

Furthermore, we suppose that the greatest eigenvalue λ_{\max}

of the matrix $\Lambda(t) = \begin{pmatrix} \sigma_1(t)Q_1 p_\alpha^\lambda & \sigma_1(t)Q_1 p_\alpha^\lambda \\ \sigma_2(t)Q_2 p_\beta^\gamma & \sigma_2(t)Q_2 p_\beta^\gamma \end{pmatrix}$, does

not exceed unity, i.e. satisfies the inequality (1.7)

Lemma 3.1: Let the vector functions $f(t, \varepsilon, u)$ and $g(t, \mu, v)$ are defined and continuous on the domain (3.1) and satisfy (1.3), then the inequality

$$\left(\begin{aligned} \|F_1(t, x_0, y_0)\| \\ \|G_1(t, x_0, y_0)\| \end{aligned} \right) \leq \begin{pmatrix} \sigma_1(t)Q_1 M \\ \sigma_2(t)Q_2 N \end{pmatrix}, \tag{3.3}$$

holds for $0 \leq t \leq T$, $\sigma_1(t) \leq T$, and $\sigma_2(t) \leq T$ where

$$\begin{aligned} F_1(t, x_0, y_0) &= \int_0^t e^{A(t-\tau)}(f(\tau, \varepsilon, u) \\ &\quad - \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon, u) d\tau) d\tau \end{aligned} \tag{3.4}$$

$$\begin{aligned} G_1(t, x_0, y_0) &= \int_0^t e^{B(t-\tau)}(g(\tau, \mu, v) \\ &\quad - \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu, v) d\tau) d\tau \end{aligned} \tag{3.5}$$

$$\sigma_1(t) = \frac{t(e^{\|A\|T} - 2e^{\|A\|t} + \|E\|) + T(e^{\|A\|t} - \|E\|)}{e^{\|A\|T} - \|E\|} \tag{3.6}$$

$$\sigma_2(t) = \frac{t(e^{\|B\|T} - 2e^{\|B\|t} + \|E\|) + T(e^{\|B\|t} - \|E\|)}{e^{\|B\|T} - \|E\|} \tag{3.7}$$

Proof: From (3.4), we have

$$\begin{aligned} \|F_1(t, x_0, y_0)\| &\leq \left(\|E\| - \frac{e^{\|A\|t} - \|E\|}{e^{\|A\|T} - \|E\|} \right) \int_0^t e^{A(t-\tau)} \|f(\tau, \varepsilon, u)\| d\tau \\ &\quad + \frac{e^{\|A\|t} - \|E\|}{e^{\|A\|T} - \|E\|} \int_0^T e^{A(T-\tau)} \|f(\tau, \varepsilon, u)\| d\tau \\ &\leq \sigma_1(t)Q_1 M \end{aligned} \tag{3.8}$$

Similarly, we get

$$\|F_1(t, x_0, y_0)\| \leq \sigma_2(t)Q_2 N \tag{3.9}$$

from (3.8) and (3.9), we conclude that the inequality (3.3) holds for $0 \leq t \leq T$, $\sigma_1(t) \leq T$, and $\sigma_2(t) \leq T$.

3.1 Approximation of periodic solution

Theorem 3.2: Let $f(t, \varepsilon, u)$ and $g(t, \mu, v)$ are defined on the domain (3.1), which are continuous in t, x, y and periodic in t of period T , also satisfying the inequalities (1.3) and the conditions (3.2) and (1.7), then the sequences of function defined by

$$\begin{aligned} x_{m+1}(t, x_0, y_0) &= x_0 \\ &\quad + \int_0^t e^{A(t-\tau)}(f(\tau, \varepsilon_{m+1}, u_{m+1}) \\ &\quad - \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon_{m+1}, u_{m+1}) d\tau) d\tau, \end{aligned} \tag{3.10}$$

with $x(0) = x_0, \quad m = 0, 1, 2, \dots,$

$$\begin{aligned} y_{m+1}(t, x_0, y_0) &= y_0 \\ &\quad + \int_0^t e^{B(t-\tau)}(g(\tau, \mu_{m+1}, v_{m+1}) \\ &\quad - \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu_{m+1}, v_{m+1}) d\tau) d\tau, \end{aligned} \tag{3.11}$$

$y(0) = y_0, \quad m = 0, 1, 2, \dots$

Where $\varepsilon_{m+1} = \int_a^b \lambda(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0))d\tau$,

$$\mu_{m+1} = \int_c^d \gamma(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0))d\tau,$$

$$u_{m+1} = \int_{-\infty}^t R(t - \tau)(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0))d\tau,$$

$$v_{m+1} = \int_{-\infty}^t G(t - \tau)(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0))d\tau,$$

are periodic in t of period T , uniformly convergent as $m \rightarrow \infty$ in the domain (3.1) to the limit functions $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ respectively, which are periodic in t of period T and satisfy the system of integral equations:

$$\begin{aligned} x(t, x_0, y_0) &= x_0 \\ &\quad + \int_0^t e^{A(t-\tau)}(f(\tau, \varepsilon, u) \\ &\quad - \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon, u) d\tau) d\tau, \end{aligned} \tag{3.12}$$

and

$$\begin{aligned} y(t, x_0, y_0) &= y_0 \\ &\quad + \int_0^t e^{B(t-\tau)}(g(\tau, \mu, v) \\ &\quad - \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu, v) d\tau) d\tau, \end{aligned} \tag{3.13}$$

where $\varepsilon = \int_a^b \lambda(x(\tau, x_0, y_0) - y(\tau, x_0, y_0))d\tau$,

$$\mu = \int_c^d \gamma(x(\tau, x_0, y_0) - y(\tau, x_0, y_0))d\tau,$$

$$u = \int_{-\infty}^t R(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0))d\tau,$$

$$v = \int_{-\infty}^t G(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0))d\tau,$$

its unique solution of (1.1) and satisfy the following inequality

$$\begin{pmatrix} \|x^0(t, x_0, y_0) - x_m(t, x_0, y_0)\| \\ \|y^0(t, x_0, y_0) - y_m(t, x_0, y_0)\| \end{pmatrix} \leq \Lambda^m (E - \Lambda)^{-1} V_0 \quad (3.14)$$

Where $V_0 = \begin{pmatrix} TQ_1M \\ TQ_2N \end{pmatrix}$, E is an identity matrix.

Proof: The sequences of functions $x_m(t, x_0, y_0)$ and $y_m(t, x_0, y_0)$ are defined in (3.10) and (3.11) are defined and continuous in the domain (3.1) and periodic in t of period T. Now by lemma 3.1 and from (3.10) when m=0, we get

$$\begin{aligned} \|x_1(t, x_0, y_0) - x_0\| &\leq \sigma_1(t)Q_1M \leq TQ_1M \\ \|y_1(t, x_0, y_0) - y_0\| &\leq \sigma_2(t)Q_2N \leq TQ_2N \end{aligned}$$

and by using mathematical induction we can prove the following inequalities for $m \geq 1$.

$$\|x_m(t, x_0, y_0) - x_0\| \leq TQ_1M \quad (3.15)$$

$$\|y_m(t, x_0, y_0) - y_0\| \leq TQ_2N \quad (3.16)$$

From (3.15) and (3.16), $x_m(t, x_0, y_0) \in D_2$ and $y_m(t, x_0, y_0) \in D_3$ for all $t \in [0, T]$, $x_0 \in D_{1f}$ and $y_0 \in D_{1g}$.

Now, we prove that the sequences $\{x_m(t, x_0, y_0)\}_{m=0}^\infty$ and $\{y_m(t, x_0, y_0)\}_{m=0}^\infty$ are uniformly convergent in the domain (3.1).

By using lemma 3.1 and from (3.10) and (3.11) when m=1, we obtain:

$$\begin{aligned} \|x_2(t, x_0, y_0) - x_1(t, x_0, y_0)\| &\leq \sigma_1(t)Q_1p_\alpha^\lambda (\|x_1(t, x_0, y_0) - x_0\| \\ &\quad + \|y_1(t, x_0, y_0) - y_0\|), \end{aligned} \quad (3.17)$$

and

$$\begin{aligned} \|y_2(t, x_0, y_0) - y_1(t, x_0, y_0)\| &\leq \sigma_2(t)Q_2p_\beta^\gamma (\|x_1(t, x_0, y_0) - x_0\| \\ &\quad + \|y_1(t, x_0, y_0) - y_0\|) \end{aligned} \quad (3.18)$$

Also by using mathematical induction, we can prove the following inequalities:

$$\begin{aligned} \|x_{m+1}(t, x_0, y_0) - x_m(t, x_0, y_0)\| &\leq \sigma_1(t)Q_1p_\alpha^\lambda (\|x_m(t, x_0, y_0) \\ &\quad - x_{m-1}(t, x_0, y_0)\| \\ &\quad + \|y_m(t, x_0, y_0) - y_{m-1}(t, x_0, y_0)\|) \end{aligned} \quad (3.19)$$

$$\begin{aligned} \|y_{m+1}(t, x_0, y_0) - y_m(t, x_0, y_0)\| &\leq \sigma_2(t)Q_2p_\beta^\gamma (\|x_m(t, x_0, y_0) \\ &\quad - x_{m-1}(t, x_0, y_0)\| \\ &\quad + \|y_m(t, x_0, y_0) - y_{m-1}(t, x_0, y_0)\|) \end{aligned} \quad (3.20)$$

we rewrite (3.19) and (3.20) with vectors form and on the following mode:

$$V_{m+1}(t) \leq \Lambda(t)V_m(t) \quad (3.21)$$

where

$$V_{m+1} = \begin{pmatrix} \|x_{m+1}(t, x_0, y_0) - x_m(t, x_0, y_0)\| \\ \|y_{m+1}(t, x_0, y_0) - y_m(t, x_0, y_0)\| \end{pmatrix}$$

$$V_m = \begin{pmatrix} \|x_m(t, x_0, y_0) - x_{m-1}(t, x_0, y_0)\| \\ \|y_m(t, x_0, y_0) - y_{m-1}(t, x_0, y_0)\| \end{pmatrix}$$

and $\Lambda(t) = \begin{pmatrix} \sigma_1(t)Q_1p_\alpha^\lambda & \sigma_1(t)Q_1p_\alpha^\lambda \\ \sigma_2(t)Q_2p_\beta^\gamma & \sigma_2(t)Q_2p_\beta^\gamma \end{pmatrix}$.

Now, we take the maximum value for the two sides of the inequality (3.21), we get:

$$V_{m+1} \leq \Lambda V_m \quad (3.22)$$

Where $\Lambda = \max_{t \in [0, T]} \Lambda(t)$, we obtain

$$\Lambda = \begin{pmatrix} TQ_1p_\alpha^\lambda & TQ_1p_\alpha^\lambda \\ TQ_2p_\beta^\gamma & TQ_2p_\beta^\gamma \end{pmatrix}, \text{ Also (3.22), we find} \quad (3.23)$$

$V_{m+1} \leq \Lambda^m V_0$ where $V_0 = \begin{pmatrix} TQ_1M \\ TQ_2N \end{pmatrix}$ and also we get

$$\sum_{i=1}^m V_i \leq \sum_{i=1}^m \Lambda^{i-1} V_0 \quad (3.24)$$

by condition (1.7) then the sequence (3.24) is uniformly convergent that is

$$\lim_{m \rightarrow \infty} \sum_{i=1}^m \Lambda^{i-1} V_0 = \sum_{i=1}^{\infty} \Lambda^{i-1} V_0 = (E - \Lambda)^{-1} V_0 \quad (3.25)$$

Let

$$\left. \begin{aligned} \lim_{m \rightarrow \infty} x_m(t, x_0, y_0) &= x^0(t, x_0, y_0) \\ \lim_{m \rightarrow \infty} y_m(t, x_0, y_0) &= y^0(t, x_0, y_0) \end{aligned} \right\} \quad (3.26)$$

Since the sequences of function (3.10) and (3.11) are defined and continuous in the domain $(t, x, y) \in [0, T] \times D_2 \times D_3$, then the limiting functions $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ are also defined and continuous in the same domain.

Moreover, from (3.25), (3.10) and (3.11), when $m \rightarrow \infty$, the limiting functions $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ are the solution of the integral equations (3.12) and (3.13) respectively.

Finally, we have to show that $x(t, x_0, y_0)$ and $y(t, x_0, y_0)$ are unique of periodic solution of the system (1.1).

Let $x_\infty(t, x_0, y_0)$ and $y_\infty(t, x_0, y_0)$ be another solution of (1.1), i.e.

$$\begin{aligned} x_\infty(t, x_0, y_0) &= x_0 \\ &\quad + \int_0^t e^{A(t-\tau)} f(\tau, \varepsilon, u) \\ &\quad - \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon, u) d\tau \end{aligned} \quad (3.27)$$

and

$$\begin{aligned} y_\infty(t, x_0, y_0) &= y_0 \\ &\quad + \int_0^t e^{B(t-\tau)} g(\tau, \mu, v) \\ &\quad - \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu, v) d\tau \end{aligned} \quad (3.28)$$

where

$$\begin{aligned} \varepsilon &= \int_a^b \lambda(x_\infty(\tau, x_0, y_0) - y_\infty(\tau, x_0, y_0)) d\tau, \\ \mu &= \int_c^d \gamma(x_\infty(\tau, x_0, y_0) - y_\infty(\tau, x_0, y_0)) d\tau, \\ u &= \int_{-\infty}^t R(t-\tau)(x_\infty(\tau, x_0, y_0) - y_\infty(\tau, x_0, y_0)) d\tau, \\ v &= \int_{-\infty}^t G(t-\tau)(x_\infty(\tau, x_0, y_0) - y_\infty(\tau, x_0, y_0)) d\tau, \end{aligned}$$

From (3.12) and (3.13) the following inequalities holds

$$\begin{aligned} & \|x(t, x_0, y_0) - x_\infty(t, x_0, y_0)\| \\ & \leq \sigma_1(t) Q_1 p_\alpha^\lambda (\|x(t, x_0, y_0) \\ & - x_\infty(t, x_0, y_0)\| \\ & + \|y(t, x_0, y_0) \\ & - y_\infty(t, x_0, y_0)\|) \end{aligned} \quad (3.29)$$

$$\begin{aligned} & \|y(t, x_0, y_0) - y_\infty(t, x_0, y_0)\| \\ & \leq \sigma_2(t) Q_2 p_\beta^\gamma (\|x(t, x_0, y_0) \\ & - x_\infty(t, x_0, y_0)\| \\ & + \|y(t, x_0, y_0) \\ & - y_\infty(t, x_0, y_0)\|) \end{aligned} \quad (3.30)$$

Thus

$$\begin{aligned} & \left(\|x(t, x_0, y_0) - x_\infty(t, x_0, y_0)\| \right. \\ & \left. \|y(t, x_0, y_0) - y_\infty(t, x_0, y_0)\| \right) \\ & \leq A \left(\|x(t, x_0, y_0) - x_\infty(t, x_0, y_0)\| \right) \end{aligned} \quad (3.31)$$

by iteration we find that

$$\begin{aligned} & \left(\|x(t, x_0, y_0) - x_\infty(t, x_0, y_0)\| \right. \\ & \left. \|y(t, x_0, y_0) - y_\infty(t, x_0, y_0)\| \right) \\ & \leq A^m \left(\|x(t, x_0, y_0) - x_\infty(t, x_0, y_0)\| \right) \end{aligned} \quad (3.32)$$

But from the condition (1.7), we get $A^m \rightarrow 0$, when $m \rightarrow \infty$, hence we obtain that $x(t, x_0, y_0) = x_\infty(t, x_0, y_0)$ and $y(t, x_0, y_0) = y_\infty(t, x_0, y_0)$ which proves that the solution is unique..

3.2 Existence of periodic solution

The problem of existence solution of the system (1.1) is uniquely connected with the existence of zeros of the functions $\Delta_1(x_0, y_0) \in D_{1f} \times D_{1g} \rightarrow R$ and $\Delta_2(x_0, y_0) \in D_{1f} \times D_{1g} \rightarrow R$ which has the form

$$\begin{aligned} & \Delta_1(x_0, y_0) \\ & = Ax_0 + \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon, u) d\tau, \end{aligned} \quad (3.33)$$

$$\begin{aligned} & \Delta_2(x_0, y_0) \\ & = By_0 + \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu, v) d\tau, \end{aligned} \quad (3.34)$$

where $\varepsilon = \int_a^b \lambda(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau$,

$$\mu = \int_c^t \gamma(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau,$$

$$u = \int_{-\infty}^t R(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau,$$

$$v = \int_{-\infty}^t G(t - \tau)(x(\tau, x_0, y_0) - y(\tau, x_0, y_0)) d\tau,$$

Since these functions (3.33) and (3.34) are approximately determined from the sequences of functions.

$$\begin{aligned} & \Delta_{1m}(x_0, y_0) \\ & = Ax_0 \\ & + \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon_{m+1}, u_{m+1}) d\tau, \end{aligned} \quad (3.35)$$

$$\begin{aligned} & \Delta_{2m}(x_0, y_0) \\ & = By_0 \\ & + \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu_{m+1}, v_{m+1}) d\tau, \end{aligned} \quad (3.36)$$

where

$$\varepsilon_{m+1} = \int_a^b \lambda(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0)) d\tau,$$

$$\mu_{m+1} = \int_c^t \gamma(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0)) d\tau,$$

$$u_{m+1} = \int_{-\infty}^t R(t - \tau)(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0)) d\tau$$

$$v_{m+1} = \int_{-\infty}^t G(t - \tau)(x_m(\tau, x_0, y_0) - y_m(\tau, x_0, y_0)) d\tau,$$

for $m = 0, 1, 2, \dots$

Theorem 3.3: Let all assumptions and conditions of theorem 3.2 be satisfied, then the following inequalities hold

$$\begin{aligned} & \|\Delta_1(x_0, y_0) - \Delta_{1m}(x_0, y_0)\| \\ & \leq \langle (L_1 Q_1 p_\alpha^\lambda \quad L_1 Q_1 p_\alpha^\lambda), \Lambda^m (E - \Lambda)^{-1} V_0 \rangle \\ & = \omega_{1m} \end{aligned} \quad (3.37)$$

$$\begin{aligned} & \|\Delta_2(x_0, y_0) - \Delta_{2m}(x_0, y_0)\| \\ & \leq \langle (L_2 Q_2 p_\beta^\gamma \quad L_2 Q_2 p_\beta^\gamma), \Lambda^m (E - \Lambda)^{-1} V_0 \rangle \\ & = \omega_{2m} \end{aligned} \quad (3.38)$$

where

$$L_1 = \frac{\|A\|T}{e^{\|A\|T} - \|E\|}, \quad L_2 = \frac{\|B\|T}{e^{\|B\|T} - \|E\|} \quad \text{and} \quad \omega_{1m}, \omega_{2m} \text{ are positive}$$

constant satisfy for $m \geq 0$, $x_0 \in D_{1f}$, $y_0 \in D_{1g}$ and $\langle \cdot \rangle$ denotes to the non-cross product in the Euclidean space R^n .

Proof:

By equations (3.33) and (3.35) and using (3.14), we have

$$\begin{aligned} & \|\Delta_1(x_0, y_0) - \Delta_{1m}(x_0, y_0)\| \\ & \leq \frac{\|A\|}{e^{\|A\|T} - \|E\|} [\sigma_1(t) Q_1 p_\alpha^\lambda (\|x(t, x_0, y_0) \\ & - x_m(t, x_0, y_0)\| \\ & + \|y(t, x_0, y_0) - y_m(t, x_0, y_0)\|)] \\ & \leq \langle (L_1 Q_1 p_\alpha^\lambda \quad L_1 Q_1 p_\alpha^\lambda), \Lambda^m (E - \Lambda)^{-1} V_0 \rangle \\ & = \omega_{1m} \end{aligned}$$

also from the equations (3.34) and (3.36) and using (3.14), we get

$$\begin{aligned} & \|\Delta_2(x_0, y_0) - \Delta_{2m}(x_0, y_0)\| \\ & \leq \frac{\|B\|}{e^{\|B\|T} - \|E\|} [\sigma_2(t) Q_2 p_\beta^\gamma (\|x(t, x_0, y_0) \\ & - x_m(t, x_0, y_0)\| \\ & + \|y(t, x_0, y_0) - y_m(t, x_0, y_0)\|)] \\ & \leq \langle (L_2 Q_2 p_\beta^\gamma \quad L_2 Q_2 p_\beta^\gamma), \Lambda^m (E - \Lambda)^{-1} V_0 \rangle \\ & = \omega_{2m} \end{aligned}$$

Now, we prove the following theorem taking in to account that the inequalities (3.37) and (3.38) will be satisfied for all $m \geq 0$.

Theorem 3.4: Suppose that for $m \geq 0$ the sequences of functions $\Delta_{1m}(x_0, y_0)$ and $\Delta_{2m}(x_0, y_0)$, which are defined in (3.35) and (3.36) satisfy the inequalities:

$$\left. \begin{aligned} & \min_{\substack{a_1+TQ_1M \leq x_0 \leq b_1-TQ_1M \\ c_1+TQ_2N \leq y_0 \leq d_1-TQ_2N}} \Delta_{1m}(x_0, y_0) \leq -\omega_{1m} \\ & \max_{\substack{a_1+TQ_1M \leq x_0 \leq b_1-TQ_1M \\ c_1+TQ_2N \leq y_0 \leq d_1-TQ_2N}} \Delta_{1m}(x_0, y_0) \geq \omega_{1m} \end{aligned} \right\} \quad (3.39)$$

$$\left. \begin{aligned} & \min_{\substack{a_1+TQ_1M \leq x_0 \leq b_1-TQ_1M \\ c_1+TQ_2N \leq y_0 \leq d_1-TQ_2N}} \Delta_{2m}(x_0, y_0) \leq -\omega_{2m} \\ & \max_{\substack{a_1+TQ_1M \leq x_0 \leq b_1-TQ_1M \\ c_1+TQ_2N \leq y_0 \leq d_1-TQ_2N}} \Delta_{2m}(x_0, y_0) \geq \omega_{2m} \end{aligned} \right\} \quad (3.40)$$

Then the system (1.1) has a periodic solutions $x = x(t, x_0, y_0)$ and $y = y(t, x_0, y_0)$ such that $x_0 \in [a_1 + TQ_1M, b_1 - TQ_1M]$, $y_0 \in [c_1 + TQ_2N, d_1 - TQ_2N]$.

Proof: Let x_1, x_2 be any points in the interval $[a1 + TQ_1M, b1 - TQ_1M]$, and y_1, y_2 be any points in the interval $[c1 + TQ_2N, d1 - TQ_2N]$ such that

$$\left. \begin{aligned} \Delta_{1m}(x_1, y_1) &= \left. \begin{aligned} \min_{\substack{a1+TQ_1M \leq x_0 \leq b1-TQ_1M \\ c1+TQ_2N \leq y_0 \leq d1-TQ_2N}} \Delta_{1m}(x_0, y_0) \\ \Delta_{1m}(x_2, y_2) &= \max_{\substack{a1+TQ_1M \leq x_0 \leq b1-TQ_1M \\ c1+TQ_2N \leq y_0 \leq d1-TQ_2N}} \Delta_{1m}(x_0, y_0) \end{aligned} \right\} (3.41) \end{aligned}$$

$$\left. \begin{aligned} \Delta_{2m}(x_1, y_1) &= \left. \begin{aligned} \min_{\substack{a1+TQ_1M \leq x_0 \leq b1-TQ_1M \\ c1+TQ_2N \leq y_0 \leq d1-TQ_2N}} \Delta_{2m}(x_0, y_0) \\ \Delta_{2m}(x_2, y_2) &= \max_{\substack{a1+TQ_1M \leq x_0 \leq b1-TQ_1M \\ c1+TQ_2N \leq y_0 \leq d1-TQ_2N}} \Delta_{2m}(x_0, y_0) \end{aligned} \right\} (3.42) \end{aligned}$$

By using the inequalities (3.37) and (3.38), we have

$$\left. \begin{aligned} \Delta_1(x_1, y_1) &= \Delta_{1m}(x_1, y_1) + (\Delta_1(x_1, y_1) - \Delta_{1m}(x_1, y_1)) < 0 \\ \Delta_1(x_2, y_2) &= \Delta_{1m}(x_2, y_2) + (\Delta_1(x_2, y_2) - \Delta_{1m}(x_2, y_2)) > 0 \end{aligned} \right\} (3.43)$$

$$\left. \begin{aligned} \Delta_2(x_1, y_1) &= \Delta_{2m}(x_1, y_1) + (\Delta_2(x_1, y_1) - \Delta_{2m}(x_1, y_1)) < 0 \\ \Delta_2(x_2, y_2) &= \Delta_{2m}(x_2, y_2) + (\Delta_2(x_2, y_2) - \Delta_{2m}(x_2, y_2)) > 0 \end{aligned} \right\} (3.44)$$

and from the continuity of the functions $\Delta_1(x_0, y_0)$ and $\Delta_2(x_0, y_0)$ and the inequalities (3.43) and (3.44), then there exists an isolated singular point $(x^0, y^0) = (x_0, y_0)$ and $x^0 \in [x_1, x_2], y^0 \in [y_1, y_2]$ where $\Delta_1(x_0, y_0)$ and $\Delta_2(x_0, y_0)$ are equals to zero, this mean that the system (1.1) has periodic solutions $x = x(t, x_0, y_0)$ and $y = y(t, x_0, y_0)$ for $x_0 \in [a1 + TQ_1M, b1 - TQ_1M], y_0 \in [c1 + TQ_2N, d1 - TQ_2N]$.

Theorem 3.5: Suppose that $\Delta_1(x_0, y_0) \in D_{1f} \times D_{1g} \rightarrow R$ and $\Delta_2(x_0, y_0) \in D_{1f} \times D_{1g} \rightarrow R$ which has the form $\Delta_1(x_0, y_0)$

$$= Ax_0 + \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon^0, u^0) d\tau, \quad (3.45)$$

$$\begin{aligned} \Delta_2(x_0, y_0) &= By_0 + \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu^0, v^0) d\tau, \quad (3.46) \end{aligned}$$

where

$$\varepsilon^0 = \int_a^b \lambda(x^0(\tau, x_0, y_0) - y^0(\tau, x_0, y_0)) d\tau,$$

$$\mu^0 = \int_c^d \gamma(x^0(\tau, x_0, y_0) - y^0(\tau, x_0, y_0)) d\tau,$$

$$u^0 = \int_{-\infty}^t R(t-\tau)(x^0(\tau, x_0, y_0) - y^0(\tau, x_0, y_0)) d\tau,$$

$$v^0 = \int_{-\infty}^t G(t-\tau)(x^0(\tau, x_0, y_0) - y^0(\tau, x_0, y_0)) d\tau,$$

where $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ are the limit of the sequences of functions (3.10) and (3.11) respectively, then the following inequalities hold:

$$\|\Delta_1(x_0, y_0)\| \leq I_3 + L_1 Q_1 M, \quad (3.47)$$

$$\|\Delta_2(x_0, y_0)\| \leq I_4 + L_2 Q_2 N, \quad (3.48)$$

$$\begin{aligned} \|\Delta_1(x_0^1, y_0^1) - \Delta_1(x_0^2, y_0^2)\| &\leq W_4 \|x_0^1 - x_0^2\| \\ &+ W_5 \|y_0^1 - y_0^2\|, \quad (3.49) \end{aligned}$$

$$\begin{aligned} \|\Delta_2(x_0^1, y_0^1) - \Delta_2(x_0^2, y_0^2)\| &\leq W_6 \|x_0^1 - x_0^2\| \\ &+ W_7 \|y_0^1 - y_0^2\|, \quad (3.50) \end{aligned}$$

where $I_3 = \|A\| \|x_0\|, I_4 = \|B\| \|y_0\|, W_1 = (1 - TQ_1 p_\alpha^\lambda)^{-1}, W_2 = (1 - TQ_2 p_\beta^\gamma)^{-1}, W_3 = (1 - T^2 Q_1 Q_2 p_\alpha^\lambda p_\beta^\gamma W_1 W_2)^{-1}, W_4 = \|A\| + L_1 Q_1 p_\alpha^\lambda W_1 W_3 (1 + TQ_2 p_\beta^\gamma W_2)$

$$W_5 = L_1 Q_1 p_\alpha^\lambda W_2 W_3 (1 + TQ_1 p_\alpha^\lambda W_1), W_6 = L_2 Q_2 p_\beta^\gamma W_1 W_3 (1 + TQ_2 p_\beta^\gamma W_2), W_7 = \|B\| + L_2 Q_2 p_\beta^\gamma W_2 W_3 (1 + TQ_1 p_\alpha^\lambda W_1)$$

Proof: From the properties of the functions $x^0(t, x_0, y_0)$ and $y^0(t, x_0, y_0)$ were fixed in the theorem 3.2, the function $\Delta_1(x_0, y_0)$ and $\Delta_2(x_0, y_0)$ are continuous and bounded by $I_3 + L_1 Q_1 M$, and $I_4 + L_2 Q_2 N$ respectively in the domain (3.1). By using (3.45) and (3.46), we get

$$\begin{aligned} \|\Delta_1(x_0, y_0)\| &\leq \|A\| \|x_0\| \\ &+ \frac{\|A\|}{e^{\|A\|T} - E} \int_0^T \|e^{A(T-\tau)}\| \|f(\tau, \varepsilon^0, u^0)\| d\tau \\ &\leq I_3 + L_1 Q_1 M \end{aligned}$$

and

$$\begin{aligned} \|\Delta_2(x_0, y_0)\| &\leq \|B\| \|y_0\| \\ &+ \frac{\|B\|}{e^{\|B\|T} - E} \int_0^T \|e^{B(T-\tau)}\| \|g(\tau, \mu^0, v^0)\| d\tau \\ &\leq I_4 + L_2 Q_2 N \end{aligned}$$

By using (3.45) and (3.46), we obtain

$$\begin{aligned} \|\Delta_1(x_0^1, y_0^1) - \Delta_1(x_0^2, y_0^2)\| &\leq \|A\| \|x_0^1 - x_0^2\| \\ &+ L_1 Q_1 p_\alpha^\lambda (\|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ &+ \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\|), \quad (3.51) \end{aligned}$$

$$\begin{aligned} \|\Delta_2(x_0^1, y_0^1) - \Delta_2(x_0^2, y_0^2)\| &\leq \|B\| \|x_0^1 - x_0^2\| \\ &+ L_2 Q_2 p_\beta^\gamma (\|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ &+ \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\|), \quad (3.52) \end{aligned}$$

The norm $\|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\|$ and $\|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\|$ in (3.51) and (3.52), were gain from the following equations:

$$\begin{aligned} x(t, x_0^1, y_0^1) &= x_0^1 \\ &+ \int_0^t e^{A(t-\tau)} (f(\tau, \varepsilon, u)) \\ &- \frac{A}{e^{AT} - E} \int_0^T e^{A(T-\tau)} f(\tau, \varepsilon, u) d\tau, \quad (3.53) \end{aligned}$$

and

$$\begin{aligned} y(t, x_0^1, y_0^1) &= x_0^1 \\ &+ \int_0^t e^{B(t-\tau)} (g(\tau, \mu, v)) \\ &- \frac{B}{e^{BT} - E} \int_0^T e^{B(T-\tau)} g(\tau, \mu, v) d\tau, \quad (3.54) \end{aligned}$$

where

$$\varepsilon = \int_a^b \lambda(x(\tau, x_0^1, y_0^1) - y(\tau, x_0^1, y_0^1)) d\tau,$$

$$\mu = \int_c^d \gamma(x(\tau, x_0^1, y_0^1) - y(\tau, x_0^1, y_0^1)) d\tau,$$

$$u = \int_{-\infty}^t R(t-\tau)(x(\tau, x_0^1, y_0^1) - y(\tau, x_0^1, y_0^1)) d\tau,$$

$$v = \int_{-\infty}^t G(t-\tau)(x(\tau, x_0^1, y_0^1) - y(\tau, x_0^1, y_0^1)) d\tau,$$

Now, use the equation (3.53) as follows

$$\begin{aligned} & \|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ & \leq \|x_0^1 - x_0^2\| \\ & + TQ_1 p_\alpha^\lambda (\|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ & + \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\|), \end{aligned}$$

then we can write

$$\begin{aligned} & \|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ & \leq W_1 \|x_0^1 - x_0^2\| \\ & + TQ_1 p_\alpha^\lambda W_1 \|y(t, x_0^1, y_0^1) \\ & - y(t, x_0^2, y_0^2)\|, \end{aligned} \quad (3.55)$$

Also, use the equation (3.54), we get

$$\begin{aligned} & \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\| \\ & \leq \|y_0^1 - y_0^2\| \\ & + TQ_2 p_\beta^\gamma (\|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ & + \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\|), \end{aligned}$$

then we write this equation as follows

$$\begin{aligned} & \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\| \\ & \leq W_2 \|y_0^1 - y_0^2\| \\ & + TQ_2 p_\beta^\gamma W_2 \|x(t, x_0^1, y_0^1) \\ & - x(t, x_0^2, y_0^2)\|, \end{aligned} \quad (3.56)$$

Now, substitute (3.56) in (3.55), we find that

$$\begin{aligned} & \|x(t, x_0^1, y_0^1) - x(t, x_0^2, y_0^2)\| \\ & \leq W_1 W_3 \|x_0^1 - x_0^2\| \\ & + TQ_1 p_\alpha^\lambda W_1 W_2 W_3 \|y_0^1 \\ & - y_0^2\|, \end{aligned} \quad (3.57)$$

and substitute (3.55) in (3.56), we get

$$\begin{aligned} & \|y(t, x_0^1, y_0^1) - y(t, x_0^2, y_0^2)\| \\ & \leq TQ_2 p_\beta^\gamma W_1 W_2 W_3 \|x_0^1 - x_0^2\| \\ & + W_2 W_3 \|y_0^1 - y_0^2\|, \end{aligned} \quad (3.58)$$

Finally, we substitute the inequalities (3.57) and (3.58) in (3.51) we get (3.49) and substitute the inequalities (3.57) and (3.58) in (3.52), we get (3.50).

dissertation, M. Sc. Thesis, college of Education, University of Mosul).

- Apostol, M. T. (1973). *Mathematical Analysis*, 2nd edition, Institute of Technology, Addison-Wesley, California.
- Burrill, C. W. and Knudsen, J. R. (1969). *Real variable*, Holt Rinehart and Winson, Inc, USA.
- Butris, R. N., & Dawoud, S. A. Solution of Integro-differential Equation of the first order with the operators. *International Journal of Scientific Engineering and Applied Science, India, (IJSEAS)*, 1.
- Byszewski, L., & Akca, H. (1997). On a mild solution of a semilinear functional-differential evolution nonlocal problem. *International Journal of Stochastic Analysis*, 10(3), 265-271.
- Lipshutz, D., & Williams, R. J. (2015). Existence, uniqueness, and stability of slowly oscillating periodic solutions for delay differential equations with nonnegativity constraints. *SIAM Journal on Mathematical Analysis*, 47(6), 4467-4535.
- Hu, J., & Li, W. P. (2005). Theory of Ordinary Differential Equations. *Existence, Uniqueness and Stability. Publications of Department of Mathematics. The Hong Kong University of Science and Technology*.
- Korol', I. I. (2010). Numerical-analytic method for the investigation of boundary-value problems for semilinear systems of differential equations. *Nonlinear Oscillations*, 13(1), 45-56.
- Korol', I. I. (2005). On periodic solutions of one class of systems of differential equations. *Ukrainian Mathematical Journal*, 57, 583-599.
- Rama, M. M., & Rao, M. (1981). Ordinary differential equations theory and applications. *United Kingdom*.
- Samoilenko, A.M. (1966) A numerical-analytic method for investigation of periodic systems of ordinary differential equations. II, *Ukrain. Mat. Zh.* 18 (2), 50-59.
- Samoilenko, A.M. (1965). A numerical-analytic method for investigation of periodic systems of ordinary differential equations. I, *Ukrain. Mat. Zh.* 17 (4), 82-93.
- Tidke, H. L., & More, R. T. (2015). Existence and uniqueness of solution of integrodifferential equation in cone metric space. *SOP Trans. Appl. Math*, 2(1), 13-19.

REFERENCES

Aziz, M. A. (2006). *Periodic solutions for some systems of non-linear ordinary differential equations* (Doctoral

كورتيا ليكولينى:

دقن فه كولينيديا، خاندنا هبوون و تاك وجيگيرا شيكارريين خولى بو سيسته ميټ هاوكيشين جياكارى وتهواوكارى يټ نه هټلى. پشت بهستن ل سهر رټكين شلوفه كرريين ژماره يى بو خاندنا شيكارټ خولى بو هاوكيشين جياكارريين ساده ژ لايى A. Samoilenko هاتيبه پيشكيشكرن.

خلاصة البحث:

في هذا البحث، دراسة الوجود، الوحدانية والاستقرارية للحلول الدورية للأنظمة المعادلات التفاضلية التكاملية اللاخطية التي تعتمد على الطرق التحليلية A. Samoilenko العددية للدراسة للحلول الدورية للمعادلة التفاضلية العادية التي قدم من قبل.