

## COMPARISON BETWEEN THE MOST FREQUENT FUNGAL SPECIES COLONIZING GRAPEVINE BERRIES AT DIFFERENT MATURATION STAGES

ASIA ABDULHAMID SAADULLAH AND SAMIR KHALAF ABDULLAH

Dept. of Biology, Faculty of Science, University of Duhok, Kurdistan Region-Iraq

Dept. of Biology, Faculty of Science, University of Zakho, Kurdistan Region-Iraq

(Accepted for publication: June 9, 2013)

### ABSTRACT

The study was undertaken to investigate the most frequent fungal species present on the surface of grape berries at different maturation at vineyards from Duhok governorate, Kurdistan region of Iraq. The associated fungi were detected both before and after surface disinfection on Dichloran Rose Bengal Chlorophenicol agar (DRBC) medium. The most abundant genera and species found on all the three stages of maturation by decreasing orders were *Aspergillus* (*A.niger*, *A.carbonarius*, *A.tubingensis*), *Cladosporium cladosporoides*, *Penicillium* (*P.brevicompactum*, *P.citrinum*, *P.glabrum*), *Alternaria alternata* and *Aureobasidium pullulans*. This result support the theory that risk of ochratoxin A contamination can be defined based on the mycobiota present on surface of healthy grape berries.

**Keywords:** Grapes, Fungi, Iraq.

### INTRODUCTION

Grapevine (*Vitis vinifera* L.) is a crop of major economic importance used for the production of table and wine grapes, raisins, juices and vinegar. The grape cultivation area in Iraq is about 48 000 hectares with production of about 265 000 tons (FAO, 2003). Duhok province in Kurdistan Region of Iraq is a major area in Iraq for cultivation and production of grapevine.

Fungal conidia are usually present on berry skins from setting and increase in number from early version to harvest, with a peak at ripening (Serra *et al.*, 2006). Black aspergilli overwinter in soil and frequent soil cultivation can favor fungal infection in vineyard. The severity of *Aspergillus* rot is influenced by excessive irrigation prior to ripening, which causes berry splitting. Rain prior to ripening which causes berry splitting favoring *Aspergillus* infection. Berry damage caused by insects, birds or other fungal infection is the primary factor affecting disease development and ochratoxin A accumulation in berries (Cozzi *et al.*, 2006; Leong *et al.*, 2006).

Black aspergilli and other moulds are found on the surface of healthy grapes at all stages of maturation and are responsible for post-harvest decay of fresh fruits (Snowdon, 1990; Serra *et al.*, 2003, 2006). Black aspergilli are also the main source of Ochratoxin A contamination of grapes and grapevine products (Leong *et al.*, 2007).

The interior of grape berry is sterile, comprising fleshy mesocarp tissue (pulp) and seeds. The berry exocarp (skin) is the primary barrier to fungal infection, consisting of a waxy cuticle, and epidermal and sub-epidermal cells. The skin can be damaged by disease, pests and environmental conditions resulting in the splitting of the fruit (Amerine *et al.*, 1972). Grapes contain high levels of sugars and other nutrients, and although they possess ideal water activity for microbial growth, their acidic pH makes them particularly susceptible to fungal spoilage, because a large bacterial component is eliminated since most bacteria prefer near neutral pH.

Several factors influence fungal colonization of grapes. Climatic conditions were found to have a significant effect. OTA contamination of grapes and wines were found to vary from year to year even in the same vineyard (Rousseau, 2003).

The aim of the present study was to investigate the most frequent fungal species present on the surface of healthy berries at different maturation stages at vineyard from Duhok governorate.

### MATERIALS AND METHODS

Grapes (Rashmew cultivar) were collected from June to September 2010 at three developmental stages of the berries. The maturation stages corresponding to pea berry (June/early July), early version (late July/

August) and ripe berry or harvest time (late August/ September).

At each sampling time in each vineyard, 20 bunches were collected by following the two diagonals (along two crossing diagonals transects). The samples were taken to the laboratory in closed paper bags, transported in cooled boxes and analyzed in the shortest time possible, usually within 24 hours of collection.

#### **Isolation from grape fruits samples:-**

From each stage and from each sample 60 berries were randomly selected, and 30 of them were surface disinfected with sodium hypochlorite 2% for 1 min., then washed twice with sterile distilled water as described by Pitt and Hocking (1997), the other 30 berries were left without surface disinfection. Ten berries from each of surface disinfected and non-disinfected lots were put per each plate containing Dichloran Rose Bengal Chloromphenicol Agar medium (DRBC) (Fluka-Germany). The plates were incubated at 25°C for 7 days.

#### **Identification of fungi:**

Samples from soil, fresh berries and dried vine fruits were examined daily with the help of stereomicroscope for sporulating fungi. Pure colonies were established on appropriate media for identification. Majority of detected species were identified to species level based on morphological and cultural characteristics. Fungi other than the genera *Aspergillus* and *Penicillium* were identified according to the manuals of Domsch *et al.*, 1980 and Pitt and Hocking, 1997.

For identification of species in the genera *Aspergillus* and *Penicillium*, pure colonies were grown on four media according to Klich (2002) and Samson *et al.*, (2000). The media are as follows: Czapeck Yeast Extract Agar incubated for seven days at 25°C (CYA25), Czapeck Yeast Extract Agar incubated for seven days at 37°C (CYA37), Czapeck Yeast Extract Agar with 20% Sucrose incubated for seven days at 25°C (CY20S), Malt Extract Agar (MEA) incubated for seven days at 25°C.

Ingredients and preparation of the above five media were mentioned in Klich (2002), Pitt and Hocking (1997). Each medium was supplemented with 50mg / ml chlorophenicol (SDI) to suppress bacterial growth. For each culture four plates were used, two of CYA and one each of CY20S, MEA. Each plate is inoculated at the center and incubated in the dark for seven days. One CYA is incubated at 37°C. The rest are incubated at 25°C.

All species identifications were according to the keys and descriptions provided by Klich (2002), Samson *et al.*, (2004) ; Frisvad and Samson; (2004) ; Abarca *et al.*, (2004) ; Samson *et al.*, (2007) .

#### **Statistical analysis:**

The data were expressed as mean  $\pm$  standard error (mean  $\pm$  S.E) and analyzed using an analysis of variance (ANOVA) followed by multiple mean comparisons. The data obtained from this study were converted to arcsine and analyzed using SAS program and means were compared using Duncan's multiple range test (SAS Institute Inc., Gary, NC, USA, 1999).

## **RESULTS**

The most frequent genera detected from disinfected and non-disinfected berries at different maturation stages were *Alternaria*, *Aspergillus*, *Aureobasidium*, *Cladosporium* and *Penicillium*.

Table (1) and Fig. (1) showed that there is no significance difference ( $P < 0.05$ ) in colonization of *Aureobasidium* at the pea berry stage compared to other stages.


In *Alternaria*, the colonization rate showed a *significance* difference ( $P < 0.05$ ) between the three maturation stages.

In *Cladosporium*, there is a significance differences between the three maturation stages at ( $P < 0.05$ ) in non-disinfected berries, whereas, in disinfected berries, colonization rate at harvest significantly different from the other two maturation stages.

**Table 1. Comparison between genera isolated from grape at different maturation stages.**

	<i>Aureobasidium</i> disinfectant.	<i>Aureobasidium</i> non disinfectant	<i>Alternaria</i> disinfectant	<i>Aternaria</i> non disinfectant	<i>Cladosporium</i> disinfectant	<i>Cladosporium</i> non disinfectant
<b>Pea berry</b>	2.667±0.440 <sup>a</sup>	4.000±0.342 <sup>a</sup>	5.567±0.483 <sup>a</sup>	6.267±0.744 <sup>a</sup>	5.167±0.563 <sup>a</sup>	6.367±0.816 <sup>a</sup>
<b>Early verasion</b>	3.067±0.383 <sup>a</sup>	5.500±0.569 <sup>b</sup>	6.967±0.543 <sup>b</sup>	8.000±0.681 <sup>b</sup>	5.467±0.488 <sup>a</sup>	7.667±0.675 <sup>b</sup>
<b>Harvesting</b>	2.167±0.304 <sup>a</sup>	4.333±0.251 <sup>b</sup>	3.700±0.346 <sup>c</sup>	4.000±0.437 <sup>c</sup>	2.967±0.379 <sup>b</sup>	3.533±0.439 <sup>c</sup>

- Data presented as mean ± S.E.
- The same letters mean no statistical differences.
- The different letters mean statistical differences.

**Figure1. Comparison between genera isolated from grape at different maturation stages**

The most three abundant *Aspergillus* species, *A.carbonarius*, *A.niger* and *A.tubingensis* were detected from disinfected and non – disinfected grapevine berries at the three maturation stages( Table 2 and Fig. (2). Colonization rates of *A.carbonarius* in pea berry showed a significance differences at ( $P<0.05$ ) from the other two stages of maturations in both disinfected at non-disinfected berries.

In *A.niger*, pea berry and early version stages showed a significant difference from harvest stage in disinfected berries, whereas, pea berry different significantly from early version and harvest stages in non- disinfected berries.

In *A.tubingensis*, there is a significant difference at ( $P<0.05$ ) between pea berry and the two other maturation stages.

The most three dominant *Penicillium* species detected in both disinfected and non-disinfectant


berries at three maturation stages were *P.brevicompactum*, *P. citrinum* and *P. glabrum* (Table 3and Figure 3).

There is no significant differences at ( $P<0.05$ ) for colonization rate by *P.brevicompactum*, *P. citrinum* and *P. glabrum* on the disinfected berries between the three maturation stages ,whereas there is a significant differences between non-disinfectant berries colonized by *P. brevicompactum* and *P. citrinum* at the pea berry stage with both early version at harvesting .Moreover, there is no significant differences at ( $P<0.05$ ) for colonization rate between non-disinfectant berries colonized by *P.glabrum* at the pea berry stage and early verasion, whereas both stages differ significantly from harvesting stages.

**Table 2. Comparison between *Aspergillus* species during three maturation stages of grape.**

	<i>A.niger</i> disinfectant	<i>A.carbonarius</i> non disinfectant	<i>A.carbonarius</i> disinfectant	<i>A.niger</i> non disinfectant	<i>A.tubingensis</i> disinfectant	<i>A.tubingensis</i> non- disinfectant
Pea berry	3.533±0.403 <sup>a</sup>	6.333±0.728 <sup>a</sup>	3.067±0.383 <sup>a</sup>	7.000±0.650 <sup>a</sup>	3.067±0.383 <sup>a</sup>	4.000±0.342 <sup>a</sup>
Early verasion	4.133±0.728 <sup>a</sup>	7.667±0.448 <sup>a</sup> <sup>b</sup>	3.667±0.353 <sup>a</sup> <sup>b</sup>	8.067±0.383 <sup>a</sup> <sup>b</sup>	3.233±0.364 <sup>b</sup>	6.667±0.461 <sup>b</sup>
Harvesting	6.333±0.342 <sup>b</sup>	8.267±0.643 <sup>b</sup>	4.500±0.302 <sup>b</sup>	9.067±0.514 <sup>b</sup>	2.967±0.370 <sup>b</sup>	6.267±0.503 <sup>b</sup>

- Data presented as mean ± S.E.
- The same letters mean no statistical differences.
- The different letters mean statistical differences



**Figure 2 Comparisons between *Aspergillus* species during three maturation stages of grape.**

**Table 3. comparison between *Penicillium* species during three maturation stages of grape.**

	<i>P.brevicompectum</i> disinfectant	<i>P.brevicompectum</i> non-disinfectant	<i>P.citrinum</i> disinfectant	<i>P.citrinum</i> non disinfectant	<i>P.glabrum</i> disinfectant	<i>P.glabrum</i> non- disinfectant
Pea berry	2.667±0.440 <sup>a</sup>	3.067±0.383 <sup>a</sup>	2.300±0.25 9 <sup>a</sup>	2.667±0.347 a	1.833±0.209 a	2.767±0.164 a
Early verasion	3.067±0.383 <sup>a</sup>	4.400±0.306 <sup>b</sup>	2.467±0.39 8 <sup>a</sup>	3.833±0.339 b	2.167±0.235 a	3.300±0.199 a
Harvestin g	3.667±0.399 <sup>a</sup>	4.833±0.365 <sup>b</sup>	2.967±0.37 0 <sup>a</sup>	4.333±0.305 b	2.467±0.398 a	4.033±0.344 b

- Data presented as mean ± S.E.
- The same letters mean no statistical differences.
- The different letters mean statistical differences.




Figure 3. Comparisons between *Penicillium* species during three maturation stages of grape.

## DISCUSSION

The most abundant genera found on all the three different stages of maturation by decreasing order were *Aspergillus*, *Cladosporium*, *Penicillium*, *Alternaria* and *Aureobasidium*. These fungi are well known as field fungi which occur commonly in the air, plant surfaces, plant debris and soil.

Serra *et al.*, (2005) reported *Cladosporium*, *Alternaria*, *Botrytis*, *Penicillium* and *Aspergillus* as the five most abundant genera found on grape berries in Portugal. They attributed the increase in colonization rates for the active spoilage agents such as *Aspergillus* and *Penicillium*, because conditions are more favorable for fungal invasion at harvest time, when more damage to the berries is likely to occur. The majority of *Aspergillus* species belong to black aspergilli (*Aspergillus* section *Nigri*). These include *A. carbonarius*, *A. niger* and *A. tubingensis*. Previous reports from different parts of the world, including Argentina and Brazil (De Rocha *et al.*, 2002; Magnoli *et al.*, 2003), and Europe (Serra *et al.*, 2005, 2006) supported our finding for predominance of black aspergilli on grape vine fruits.

Species of *Penicillium* showed high frequency of occurrence on different maturation stages were *P. brevicompactum*, *P. citrinum* and *P. glabrum*. The most frequent species of *Penicillium* reported on Portugies grapevine berries by Serra *et al.*, (2005) were *P. brevicompactum*, *P. thomi* and *P. glabrum*.

It is interesting to note that *Botrytis cinerea*, which is responsible for grey rot in grapes bunches, was not detected in our study. This is may be attributed to the dry and hot weather during grape vines berries maturation in our region.

*Aureobasidium pullulans* is anamorphic yeast – like fungus was also isolated from berries at different maturation stages with high colonization rate. The fungus is well known as a common inhabitant of the phylloplane of several plants. Recent studies indicated that some strains of *A. pullulans* used as biocontrol agents by preventing *A. carbonarius* infection to berries. Moreover, berries pretreated with the biocontrol agent and infected with OTA as compared to untreated infected control berries, the strains showed ability to degrade OTA to ochratoxin A in berries which is less toxic (De Felice *et al.*, 2008).

In conclusion, the presence of toxigenic fungi particularly black aspergilla on the skin of berries at different maturation stages will favor toxin production. The skin can be damaged by pests and environmental conditions resulting in the splitting of the fruit (Amerine *et al.* 1972). Grapes contain high level of sugars and other nutrients; their acidic PH make them potentially susceptible for fungal spoilage and ochratoxin A contamination. Among black aspergilli, *A. carbonarius* is the most important as ochratoxin A producing isolates, followed by *A. niger* aggregate (Varga *et al.* 2004).

## REFERENCES

- Abarca, M.L.; Accensi, F.; Cano, J. and Cabanes, F.J. (2004). Taxonomy and significance of black aspergilla. Antonio Van Leeuwenhoek 86, 33-49.
- Amerine, M. A., Berg, H. W. and Cruess, W. V. (1972). Technology of Wine Making. 3rd edition. The AVI Publishing Company Inc. Westport, Connecticut, USA
- Cozzi, G., Pascale, M., Perrone, G., Visconti, A. and Logrieco, A., (2006). Effect of *Lobesia botrana* damages on black aspergilli rot and ochratoxin A content in grapes. *Int. J. Food Microbiol.* 111, 10-17.
- De Felice, D.V., Solfrizzo, M., De Curtis, F., Lima, G., Visconti, A. and Castoria, R. (2008). Strains of *Aureobasidium pullulans* can lower ochratoxin A contamination in wine grapes. *Phytopathology*, 98: 1261-1270.
- De Rocha Rosa, Palacios, V., Combina, M., Fraga, M.E., De, O.R. and Magnoli, C. E. (2002). Potential ochratoxin A producers from wine grapes in Argentina and Brazil. *Food addit. Contam.* 19, 408-414.
- Domsch, K. H., Gams, W. and Anderson, T. H. (1980). Compendium of soil fungi. Academic Press, London.
- F.A.O (2003). The (UN) Bulletin of statistics, Vo.14, No.2.
- Frisvad, J. C. and Samson, R. A. (2004). Polyphasic taxonomy of *Penicillium* subgenus *Penicillium*. A guide to identification of food and air-borne terverticillate *Penicillia* and their mycotoxins. *Studies in Mycology*, 49, 1- 173.
- Frisvad, J.C., Frank, J.M., Houbraken, J.A.M.P., Kuijpers, A.F.A. and Samson, R.A., (2004). New ochratoxin A producing species of *Aspergillus* section *Circumdati*. *Stud. Mycol.*, 50, 23-43.
- Klich M.A. (2002). Identification of common *Aspergillus* species, CBS, Utrescht, The Netherlands pp.116.
- Leong, S. L., Hocking, A. D. Pitt J.I. Kazi B.A., Emme R.W. and Scott, E. S. (2006). Black *Aspergillus* species in Australian vineyards: From soil to ochratoxin A in wine In :Advances in Food Mycology, Hocking, A.D., Pitt, J.I., Samson R.A. and Throne U. (eds). *Springer Science*, USA, pp 153-171.
- Leong, S. L., Hocking, A. D., and Scott, E. S. (2007). *Aspergillus* species producing ochratoxin A: isolation from vineyard soil and infection of Semillon bunches in Australia *J. Appl. Microbiol.* 124-133.
- Magnoli, C., Violante, M., Combina, M., Palacio, G., and Dalcero, A. (2003). Mycoflora and ochratoxin-producing strains of *Aspergillus* section *Nigri* in wine grapes in Argentina. *Letters. Appl. Microbiol.*, 37, 179-184.
- Pitt, J. I. and Hocking, A. D. (1997). Fungal and Food Spoilage 2<sup>nd</sup> ed. London, U.K.: Blackie Academic & Professional. pp.511.
- Rousseau, J. (2003). Etude des facteurs de prevention et de reduction de la teneur en OTA dans les vine. Poster du Symposium International d' Oenologie, Bordeaux
- Samson, R. A., Hoekstra, E. S., Frisvad, J. C. and Filtenborg, O. (2000). Introduction to Food- and Air-borne Fungi 6<sup>th</sup> ed. Baarn, The Netherlands: Centraalbureau Voor Schimmelcultures.
- Samson, R. A., Houbraken, J. A. M. P., Kuijpers, A. F. A., Frank, J. M. and Frisvard, J. C. (2004). New ochratoxin A or sclerotium producing species in *Aspergillus* section *Nigri*. *Stud. Mycol.* 50, 45-61.
- Samson, R.A., Noonim, P., Meijer, M., Houbraken, J., Frisvad, J.C and Varga, J. (2007). Diagnostic tools to identify black aspergilli. *Stud. Mycol.* 59: 129-145.
- SAS (1999). SAS/STAT User's Guide, Version 8.2, 1st printing. Vol. 2. SAS Institute Inc, SAS Campus Drive, Gary, North Carolina.
- Serra, R., Abrunhosa, L., Kozakiewicz, Z. and Venâncio, A. (2003). Black *Aspergillus* species as ochratoxin A producers in Portuguese wine grapes. *Int. J. Food Microbiol.* 88, 63-68.
- Serra, R., Braga, A. and Venâncio, A. (2005). Mycotoxin-producing and other fungi isolated from grapes for wine production, with particular emphasis on ochratoxin A. *Res. Microbiol.* 156, 515-521.
- Serra, R., Mendonca, C., Venancio, A. (2006). Ochratoxin A occurrence and formation in Portuguese wine grapes at various stages of maturation. *Int. J. Food Microbiol.* 111, S35-S39.
- Snowdon, A. L. (1990). Colour Atlas of post-harvest diseases and disorders of fruits and vegetables. Wolfe Scientific, London. pp.256.
- Varga, J., Juhasz, A., Kevei, F., and Kozakiewicz, Z. (2004). Molecular diversity of agriculturally important *Aspergillus* species. *Eur. J. plant pathol.*, 110, 627-640.

#### الخلاصة

اجريت هذه الدرسة لمعرفة اهم الانواع الفطرية المتواجدة علي سطح ثمار العنب لمراحل النضج المختلفة في محافظة دهوك اقليم كوردستان العراق عزلت الفطريات قبل و بعد تعقيم السطح التمار على وسط DRBC الاجناس والانواع الساعدة في جميع المراحل الثلاثة للنضج هي

*Aspergillus* (*A.niger*, *A.carbonarius*, *A.tubingensis*), *Cladosporium cladosporoides*, *Penicillium* (*P.brevicompactum* , *P.citrinum*, *P.glabrum*), *Alternaria alternata* and *Aureobasidium pullalans* دعمت هذه النتيجة النظرية القائلة بان التلوث بالسم الفطري ochratoxin A يحدد اعتمادا علي الانواع الفطرية الموجودة على سطح التمار السليمة.

#### بوغته

ئهف فهكولينه هاتنه كرن دا فهكولين لسهر هه بونا كومين كه رويالسه ر به رههمي تري د قوناغين جوره جور ديبكغه هشتي دا ل پاريزگه ها دهوكي، هيريما كوردستانا عراقي، فافارتنا كه روا هاته كرن بهري و پشتي تافيكرونا ره خوريت نه وان ز پشتي هنگي هاته دانان زناف نامانين چاندني كو بيكهاتيه ژ DRBC نهو جوره كه زين به ربه ثاف د هه رسي قوناغين بيكغه هشتيدا فه دگرين بو *Aspergillus* (*A.niger*, *A.carbonarius*, *A.tubingensis*), *Cladosporium cladosporoides*, *Penicillium* (*P.brevicompactum* ,*P.citrinum*,*P.glabrum*), *Alternaria alternata* and *Aureobasidium pullalans* نه نجاما دياركري نه نجاما ديارى نهو تيوروا ديبثيت پيسبونا تري بژهره كه روا ochratoxin A دهيته دهست نيشان كرن ل جورين كومه كه رويين ل سه ر تري ساخله م .